arXiv:1406.4122v1 [math-ph] 14 Jun 2014

(Pseudo) Generalized Kaluza-Klein G-Spaces and Einstein Equations

C. M. Arcuş and E. Peyghan

June 18, 2014

Abstract

Introducing the Lie algebroid generalized tangent bundle of a Kaluza-Klein bundle, we develop the theory of general distinguished linear connections for this space. In particular, using the Lie algebroid generalized tangent bundle of the Kaluza-Klein vector bundle, we present the (g,h)-lift of a curve on the base M and we characterize the horizontal and vertical parallelism of the (g,h)-lift of accelerations with respect to a distinguished linear (ρ,η)-connection. Moreover, we study the torsion, curvature and Ricci tensor field associated to a distinguished linear (ρ,η)-connection and we obtain the identities of Cartan and Bianchi type in the general framework of the Lie algebroid generalized tangent bundle of a Kaluza-Klein bundle. Finally, we introduce the theory of (pseudo) generalized Kaluza-Klein G-spaces and we develop the Einstein equations in this general framework.

1 Introduction

Recently, Lie algebroids are important issues in physics and mechanics since the extension of Lagrangian and Hamiltonian systems to their entity [11, 13, 19, 27] and catching the poisson structure [20]. Then, Arcuş introduced generalized Lie algebroids as the extension of Lie algebroids and he studied geometrical and physical concepts for these spaces [1, 2, 3, 4]. Indeed a generalized Lie algebroid is a extension of Lie algebroid from one base manifold to a pair of diffeomorphic base manifolds.

The Einstein theory of general relativity and Maxwell theory of electromagnetism was independent developed in the world of physicists. In [12] Kaluza proposed to unify these two theories using a five-dimensional manifold. Kaluza's achievement was possible if the components of the pseudo-Riemannian metric on the five-dimensional manifold does not depend on the fifth coordinate (cylinder condition). Then, Klein [14] added the condition of compactification, namely the space is closed by a very small circle in the direction of the fifth dimension. Therefore, the Kaluza-Klein theory emerged to unify of the Einstein theory of general gravity and Maxwell theory of electromagnetism.

In [10], Einstein and Bergmann proposed a first generalization of Kaluza-Klein theory using a pseudo-Riemannian metric such that its components are periodic in the fifth coordinate. So, the cylinder condition was partially satisfied and, for the firs time, a covariant derivative of a vector bundle over the five-dimensional manifold was introduced.

A well known generalization of the five-dimensional Kaluza-Klein theory is the so called the space time matter (STM) theory. Cosmological solutions in which both the cylinder condition and the compactification condition were removed are presented in [9, 15, 17, 18]. An excelent survey on STM theory is presented in the paper of Overduin and Wesson [16].

The idea to construct exact solutions with Lie and Clifford algebroid symmetries in modified and extra dimension gravity and matter field theories was elaborated originally in a series of preprints by S. Vacaru [21, 22, 23]; see further developments and reviews of results on nonholonomic algebroids and Einstein-Direact structures, Finsler-Lagrange-Hamilton algebroid spaces and geometric flows on Lie algebroid in [24, 25, 26].

Recently, Bejancu developed a new point of view on a general Kaluza-Klein theory using the product manifold $\overline{M} = M \times K$, where M is a four-dimensional manifold and K is a one-dimensional manifold [5, 6, 7]. The tangent bundle of $\overline{M}$ is the space used to develop the theory as a direct sum between the horizontal distribution $H\overline{M}$ and vertical distribution $V\overline{M}$. The novelty is determined by the Riemannian horizontal connection which plays the same role as Levi-Civita linear connection on the space time manifold M for the classical Kaluza-Klein theory. Using this linear connection, which is not distinguished connection, Bejancu introduced the Einstein gravitational tensor field and he write the field equations on $\left(\overline{M}, \overline{g}\right)$. A new method for the study of general higher dimension Kaluza-Klein theory is presented by Bejancu in [8].

We remark that Bejancu used the usual Lie algebroid tangent bundle. The purpose of this paper is to develop a general method to study of the Kaluza-Klein theory using the generalized connection theory presented by Arcuş in [1, 2, 3, 4].

This paper is arranged as follows. In Sec. 3, using a (generalized) Lie algebroid presented in Sec. 2, we introduce the Lie algebroid generalized tangent bundle of a Kaluza-Klein bundle and we study the connections on a Kaluza-Klein bundle. Moreover, we develop the theory of general distinguished linear connections for the Lie algebroid generalized tangent bundle. In Sec. 4, using the Lie algebroid generalized tangent bundle of the Kaluza-Klein vector bundle, we present the (g, h)-lift of a curve on the base M. A characterization of horizontality (respectively, verticality) parallelism of the (g, h)-lift of accelerations with respect to a distinguished linear (ρ, η)-connection is presented. Sec. 5 is dedicated to study the torsion and curvature of a distinguished linear (ρ, η)-connection. Moreover, the Ricci tensor field associated to a distinguished linear (ρ, η)-connection is introduced and identities of Cartan and Bianchi type in the general framework of the Lie algebroid generalized tangent bundle of a Kaluza-Klein bundle are presented. Finally, in Sec. 6, we present the theory of (pseudo) generalized Kaluza-Klein G-spaces. A lot of examples of Kaluza-Klein

G-spaces are presented and the Einstein equations are developed in this general framework. In particular, using the identities morphisms, the usual Kaluza-Klein theory used by Bejancu is obtained, but the difference is that the metrical linear connection used in our paper is distinguished connection.

2 Preliminaries

Let (F,ν,N) be a vector bundle, $\Gamma(F,\nu,N)$ be the set of the sections of it and $\mathcal{F}(N)$ be the smooth real-valued functions on N. Then $(\Gamma(F,\nu,N),+,\cdot)$ is a $\mathcal{F}(N)$-module. If (φ,φ_0) is a morphism from (F,ν,N) to (F',ν',N') such that φ_0 is a isomorphism from N to N', then using the operation

$$\begin{array}{ccc} \mathcal{F}(N)\times\Gamma(F',\nu',N') & \stackrel{\cdot}{\longrightarrow} & \Gamma(F',\nu',N'), \\ (f,u') & \longmapsto & f\circ\varphi_0^{-1}\cdot u', \end{array}$$

it results that $(\Gamma(F',\nu',N'),+,\cdot)$ is a $\mathcal{F}(N)$-module and we obtain the modules morphism

$$\begin{array}{ccc} \Gamma(F,\nu,N) & \xrightarrow{\Gamma(\varphi,\varphi_0)} & \Gamma(F',\nu',N'), \\ u & \longmapsto & \Gamma(\varphi,\varphi_0)\,u, \end{array}$$

defined by

$$\Gamma(\varphi,\varphi_0)\,u(y)=\varphi\left(u_{\varphi_0^{-1}(y)}\right)=\left(\varphi\circ u\circ\varphi_0^{-1}\right)(y),$$

for any $y\in N'$.

Definition 1 *A generalized Lie algebroid is a vector bundle (F,ν,N) given by the diagrams:*

$$\begin{array}{ccccc} \left(F,[,]_{F,h}\right) & \xrightarrow{\rho} & (TM,[,]_{TM}) & \xrightarrow{Th} & (TN,[,]_{TN}) \\ \downarrow\nu & & \downarrow\tau_M & & \downarrow\tau_N \\ N & \xrightarrow{\eta} & M & \xrightarrow{h} & N \end{array} \tag{1}$$

where h and η are arbitrary isomorphisms, (ρ,η) is a vector bundles morphism from (F,ν,N) to (TM,τ_M,M) and

$$\begin{array}{ccc} \Gamma(F,\nu,N)\times\Gamma(F,\nu,N) & \xrightarrow{[,]_{F,h}} & \Gamma(F,\nu,N), \\ (u,v) & \longmapsto & [u,v]_{F,h}, \end{array}$$

is an operation satisfies in

$$[u,f\cdot v]_{F,h}=f\,[u,v]_{F,h}+\Gamma(Th\circ\rho,h\circ\eta)(u)\,f\cdot v,\quad\forall f\in\mathcal{F}(N),$$

such that

1) The 4-tuple $(\Gamma(F,\nu,N),+,\cdot,[,]_{F,h})$ is a Lie $\mathcal{F}(N)$-algebra,

2) The modules morphism $\Gamma(Th\circ\rho,h\circ\eta)$ is a Lie algebras morphism from $(\Gamma(F,\nu,N),+,\cdot,[,]_{F,h})$ to $(\Gamma(TN,\tau_N,N),+,\cdot,[,]_{TN})$.

We denote by $\Big((F,\nu,N),[,]_{F,h},(\rho,\eta)\Big)$ the generalized Lie algebroid defined in the above. Moreover, the couple $\Big([,]_{F,h},(\rho,\eta)\Big)$ is called the *generalized Lie algebroid structure.*

A morphism from $\Big((F,\nu,N),[,]_{F,h},(\rho,\eta)\Big)$ to $\Big((F',\nu',N'),[,]_{F',h'},(\rho',\eta')\Big)$ is a morphism (φ,φ_0) from (F,ν,N) to (F',ν',N') such that φ_0 is an isomorphism from N to N', and the modules morphism $\Gamma(\varphi,\varphi_0)$ is a Lie algebras morphism from $\Big(\Gamma\left(F,\nu,N\right),+,\cdot,[,]_{F,h}\Big)$ to $\Big(\Gamma\left(F',\nu',N'\right),+,\cdot,[,]_{F',h'}\Big)$. Thus, we can discuss about the category of generalized Lie algebroids.

Remark 2 *In the particular case, $(\eta,h)=(Id_M,Id_M)$, we obtain the definition of Lie algebroid.*

If we take local coordinates (x^i) and $(\chi^{\tilde{\imath}})$ on open sets $U\subset M$ and $V\subset N$, respectively, then we have the corresponding local coordinates (x^i,y^i) and $(\chi^{\tilde{\imath}},z^{\tilde{\imath}})$ on TM and TN, respectively, where $i,\tilde{\imath}\in 1,\ldots,m$. Moreover, a local basis $\{t_\alpha\}$ of the sections of $\nu^{-1}(V)\to V$ generates local coordinates $(\chi^{\tilde{\imath}},z^\alpha)$ on F, where $\alpha\in 1,\ldots,p$. If we consider the another local coordinates $(x^{i'}(x^i),y^{i'}(x^i,y^i))$, $(\chi^{\tilde{\imath}'}(\chi^{\tilde{\imath}}),z^{\tilde{\imath}'}(\chi^{\tilde{\imath}},z^{\tilde{\imath}}))$ and $(\chi^{\tilde{\imath}'}(\chi^{\tilde{\imath}}),z^{\alpha'}(\chi^{\tilde{\imath}},z^\alpha))$ on TM, TN and F, respectively, then we have the following corresponding changes of coordinates

$$z^{\alpha'}=\Lambda^{\alpha'}_{\alpha}z^{\alpha},\quad y^{i'}=\frac{\partial x^{i'}}{\partial x^i}y^i,\quad z^{\tilde{\imath}\prime}=\frac{\partial\chi^{\tilde{\imath}\prime}}{\partial\chi^{\tilde{\imath}}}z^{\tilde{\imath}}. \tag{2}$$

We assume that $(\theta,\mu)=(Th\circ\rho,h\circ\eta)$. If $z^\alpha t_\alpha$ is a section of (F,ν,N), then for any $f\in\mathcal{F}(N)$ and $\varkappa\in N$ we have

$$\begin{aligned}\Gamma\left(Th\circ\rho,h\circ\eta\right)\left(z^\alpha t_\alpha\right)f\left(h\circ\eta\left(\varkappa\right)\right)&=\left(\theta^{\tilde{\imath}}_\alpha z^\alpha\tfrac{\partial f}{\partial\varkappa^{\tilde{\imath}}}\right)\left(h\circ\eta\left(\varkappa\right)\right)\\&=\left(\left(\rho^i_\alpha\circ h\right)\left(z^\alpha\circ h\right)\tfrac{\partial f\circ h}{\partial x^i}\right)\left(\eta\left(\varkappa\right)\right),\end{aligned}$$

where ρ^i_α are local functions on N [1]. Now, we put $[t_\alpha,t_\beta]_{F,h}=L^\gamma_{\alpha\beta}t_\gamma$, where $L^\gamma_{\alpha\beta}$ are local functions on N and $\alpha,\beta,\gamma\in 1,\ldots,p$. It is easy to see that $L^\gamma_{\alpha\beta}=-L^\gamma_{\beta\alpha}$. Moreover, the condition (2) of *Definition 1* implies that

$$\left(L^\gamma_{\alpha\beta}\circ h\right)\left(\rho^k_\gamma\circ h\right)=\left(\rho^i_\alpha\circ h\right)\frac{\partial\left(\rho^k_\beta\circ h\right)}{\partial x^i}-\left(\rho^j_\beta\circ h\right)\frac{\partial\left(\rho^k_\alpha\circ h\right)}{\partial x^j}.$$

The local functions ρ^i_α, $L^\gamma_{\alpha\beta}$ introduced in the above are called the *structure functions* of the generalized Lie algebroid $\Big((F,\nu,N),[,]_{F,h},(\rho,\eta)\Big)$. Under the change of coordinates (2), ρ^i_α and $\theta^{\tilde{\imath}}_\alpha$ satisfy in the following transformation rules

$$\rho^{i'}_{\alpha'}=\Lambda^\alpha_{\alpha'}\rho^i_\alpha\frac{\partial x^{i'}}{\partial x^i},\quad\theta^{\tilde{\imath}'}_{\alpha'}=\Lambda^\alpha_{\alpha'}\theta^{\tilde{\imath}}_\alpha\frac{\partial\varkappa^{\tilde{\imath}'}}{\partial\varkappa^{\tilde{\imath}}},$$

where $\|\Lambda_{\alpha'}^{\alpha}\| = \left\|\Lambda_{\alpha}^{\alpha'}\right\|^{-1}$. Also, it is known that the following relation is hold between ρ_{α}^{i} and $\theta_{\alpha}^{\tilde{i}}$

$$\rho_{\alpha}^{i} \circ h \frac{\partial f \circ h}{\partial x^{i}} = \left(\theta_{\alpha}^{\tilde{i}} \frac{\partial f}{\partial \varkappa^{\tilde{i}}}\right) \circ h, \quad \forall f \in \mathcal{F}(N).$$

3 The Lie algebroid generalized tangent bundle

Let M be a 4-dimensional manifold and K be a 1-dimensional manifold. Considering $E = M \times K$, we introduce the Kaluza-Klein bundle (E, π, M), where $\pi : E \longrightarrow M$ is the projection map on the first factor. Let (x^i), $i \in \{1, 2, 3, 4\}$, be a coordinate system on M. Then we can consider a coordinate system (x^i, y°) on E, where y° is the fibre coordinate. Let

$$(x^i, y^\circ) \longrightarrow (x^{i'}(x^i), y^{\circ\prime}(x^i, y^\circ)),$$

be a change of coordinates on (E, π, M). Then the coordinate y° change to $y^{\circ\prime}$ according to the rule:

$$y^{\circ\prime} = \frac{\partial y^{\circ\prime}}{\partial y^\circ} y^\circ.$$

Now, let $\left((F, \nu, N), [,]_{F,h}, (\rho, \eta)\right)$ be a generalized Lie algebroid. Using the diagram

$$\begin{array}{ccc} E & & \left(F, [,]_{F,h}, (\rho, \eta)\right) \\ \pi \downarrow & & \downarrow \nu \\ M & \xrightarrow{\quad h \quad} & N \end{array} \tag{3}$$

we have the pull-back bundle $(\pi^*(h^*F), \pi^*(h^*\nu), E)$. Here we consider the vector bundles morphism $\left(\overset{\pi^*(h^*F)}{\rho}, Id_E\right)$ from $(\pi^*(h^*F), \pi^*(h^*\nu), E)$ to (TE, τ_E, E), where

$$\begin{array}{ccc} \pi^*(h^*F) & \overset{\pi^*(h^*F)}{\rho} & TE, \\ Z^\alpha T_\alpha(u_x) & \longmapsto & \left(Z^\alpha \cdot \rho_\alpha^i \circ h \circ \pi\right) \dfrac{\partial}{\partial x^i}(u_x), \end{array}$$

and $\{T_\alpha\}_{\alpha=1}^{p}$ be a basis of the sections of the pull-back bundle

$$(\pi^*(h^*F), \pi^*(h^*\nu), E).$$

Using the operation

$$\Gamma(\pi^*(h^*F), \pi^*(h^*\nu), E)^2 \xrightarrow{[,]_{\pi^*(h^*F)}} \Gamma(\pi^*(h^*F), \pi^*(h^*\nu), E),$$

defined by

$$\begin{aligned} [T_\alpha, T_\beta]_{\pi^*(h^*F)} &= L_{\alpha\beta}^{\gamma} \circ h \circ \pi \cdot T_\gamma, \\ [T_\alpha, fT_\beta]_{\pi^*(h^*F)} &= fL_{\alpha\beta}^{\gamma} \circ h \circ \pi T_\gamma + \rho_\alpha^i \circ h \circ \pi \frac{\partial f}{\partial x^i} T_\beta, \\ [fT_\alpha, T_\beta]_{\pi^*(h^*F)} &= -[T_\beta, fT_\alpha]_{\pi^*(h^*F)}, \end{aligned}$$

for any $f \in \mathcal{F}(E)$, it results that

$$\left(\left(\pi^* \left(h^* F \right), \pi^* \left(h^* \nu \right), E \right), [,]_{\pi^*(h^*F)}, \left(\overset{\pi^*(h^*F)}{\rho}, Id_E \right) \right),$$

is a Lie algebroid which is called the *pull-back Lie algebroid* of the generalized Lie algebroid $\left((F, \nu, N), [,]_{F,h}, (\rho, \eta) \right)$.

If $z = z^\alpha t_\alpha \in \Gamma(F, \nu, N)$, then we obtain the section

$$Z = \left(z^\alpha \circ h \circ \pi \right) T_\alpha \in \Gamma \left(\pi^* \left(h^* F \right), \pi^* \left(h^* \nu \right), E \right),$$

such that $Z(u_x) = z(h(x))$, for any $u_x \in \pi^{-1}\left(U \cap h^{-1} V \right)$.

Now we consider the vector bundle $\left(\pi^* \left(h^* F \right) \oplus TE, \overset{\oplus}{\pi}, E \right)$. Let $\left(\partial_i, \dot{\partial}_\circ \right)$ be the base sections for the Lie $\mathcal{F}(E)$-algebra

$$\left(\Gamma(TE, \tau_E, E), +, \cdot, [,]_{TE} \right),$$

where $\partial_i := \frac{\partial}{\partial x^i}$ and $\dot{\partial}_\circ := \frac{\partial}{\partial y^\circ}$. Setting

$$\tilde{\partial}_\alpha := T_\alpha \oplus (\rho_\alpha^i \circ h \circ \pi) \partial_i, \quad \dot{\tilde{\partial}}_\circ := 0_{\pi^*(h^*F)} \oplus \dot{\partial}_\circ,$$

one can deduce that

$$Z^\alpha \tilde{\partial}_\alpha + Y^\circ \dot{\tilde{\partial}}_\circ = Z^\alpha T_\alpha \oplus \left(Z^\alpha \left(\rho_\alpha^i \circ h \circ \pi \right) \partial_i + Y \dot{\partial}_\circ \right),$$

is a section of $\left(\pi^* \left(h^* F \right) \oplus TE, \overset{\oplus}{\pi}, E \right)$, where $Z^\alpha T_\alpha$ and $Y^\circ \dot{\partial}_\circ$ are sections of $\left(\pi^* \left(h^* F \right), \pi^* \left(h^* F \right), E \right)$ and (VTE, τ_E, E), respectively. Moreover, it is easy to see that the sections $\tilde{\partial}_1, \ldots, \tilde{\partial}_p, \dot{\tilde{\partial}}_\circ$ are linearly independent. Therefore we can consider the vector subbundle $((\rho, \eta) TE, (\rho, \eta) \tau_E, E)$ of the vector bundle $\left(\pi^* \left(h^* F \right) \oplus TE, \overset{\oplus}{\pi}, E \right)$, for which the $\mathcal{F}(E)$-module of sections is the $\mathcal{F}(E)$-submodule of $\left(\Gamma \left(\pi^* \left(h^* F \right) \oplus TE, \overset{\oplus}{\pi}, E \right), +, \cdot \right)$ generated by the set of sections $\left(\tilde{\partial}_\alpha, \dot{\tilde{\partial}}_\circ \right)$. The vector bundle $((\rho, \eta) TE, (\rho, \eta) \tau_E, E)$ is called the *generalized tangent bundle* and the base sections $\left(\tilde{\partial}_\alpha, \dot{\tilde{\partial}}_\circ \right)$ is called the *natural* (ρ, η)-base.

The matrix of coordinate transformation on $((\rho, \eta) TE, (\rho, \eta) \tau_E, E)$ at a change of fibred charts is

$$\begin{pmatrix} \Lambda_\alpha^{\acute{\alpha}} \circ h \circ \pi & 0 \\ \left(\rho_\alpha^i \circ h \circ \pi \right) \dfrac{\partial y^{\circ\prime}}{\partial x^i} & \dfrac{\partial y^{\circ\prime}}{\partial y^\circ} \end{pmatrix}.$$

Theorem 3 *The vector bundle $((\rho, \eta) TE, (\rho, \eta) \tau_E, E)$ has a Lie algebroid structure.*

Proof. We consider the vector bundles morphism $(\tilde{\rho}, Id_E)$ from $((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E)$ to (TE, τ_E, E), where

$$(\rho,\eta)\,TE \xrightarrow{\tilde{\rho}} TE$$
$$\left(Z^\alpha \tilde{\partial}_\alpha + Y^\circ \dot{\tilde{\partial}}_\circ\right)(u_x) \longmapsto \left(Z^\alpha \left(\rho^i_\alpha \circ h \circ \pi\right)\partial_i + Y^\circ \dot{\partial}_\circ\right)(u_x).$$

Also, we define the bracket

$$\Gamma\left((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E\right)^2 \xrightarrow{[,]_{(\rho,\eta)TE}} \Gamma\left((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E\right),$$

by

$$\left[Z_1^\alpha \tilde{\partial}_\alpha + Y_1^\circ \dot{\tilde{\partial}}_\circ, Z_2^\beta \tilde{\partial}_\beta + Y_2^\circ \dot{\tilde{\partial}}_\circ\right]_{(\rho,\eta)TE} = \left[Z_1^\alpha T_\alpha, Z_2^\beta T_\beta\right]_{\pi^*(h^*F)}$$
$$\oplus \left[Z_1^\alpha \left(\rho^i_\alpha \circ h \circ \pi\right)\partial_i + Y_1^\circ \dot{\partial}_\circ,\ Z_2^\beta \left(\rho^j_\beta \circ h \circ \pi\right)\partial_j + Y_2^\circ \dot{\partial}_\circ\right]_{TE},$$

for any sections $Z_1^\alpha \tilde{\partial}_\alpha + Y_1^\circ \dot{\tilde{\partial}}_\circ$ and $Z_2^\beta \tilde{\partial}_\beta + Y_2^\circ \dot{\tilde{\partial}}_\circ$ of $((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E)$. It is easy to check that $\left([,]_{(\rho,\eta)TE}, (\tilde{\rho}, Id_E)\right)$ is a Lie algebroid structure for the vector bundle $((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E)$. ■

Remark 4 *In particular, if $h = Id_M$, then the Lie algebroid*

$$\left(((Id_{TM}, Id_M)\,TE, (Id_{TM}, Id_M)\,\tau_E, E), [,]_{(Id_{TM}, Id_M)TE}, \left(\widetilde{Id_{TM}}, Id_E\right)\right),$$

is isomorphic with the usual Lie algebroid

$$\left((TE, \tau_E, E), [,]_{TE}, (Id_{TE}, Id_E)\right).$$

This is a reason for which the Lie algebroid

$$\left(((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E), [,]_{(\rho,\eta)TE}, (\tilde{\rho}, Id_E)\right),$$

is called the Lie algebroid generalized tangent bundle.

3.1 (ρ,η)-connections

We consider the vector bundles morphism $((\rho,\eta)\,\pi!, Id_E)$ given by the commutative diagram

$$\begin{array}{ccc} (\rho,\eta)\,TE & \xrightarrow{(\rho,\eta)\pi!} & \pi^*\left(h^*F\right) \\ (\rho,\eta)\,\tau_E \downarrow & & \downarrow \pi^*\left(h^*\nu\right) \\ E & \xrightarrow{Id_E} & E \end{array}$$

defined by

$$(\rho,\eta)\,\pi!\left(\left(Z^\alpha \tilde{\partial}_\alpha + Y^\circ \dot{\tilde{\partial}}_\circ\right)(u_x)\right) = \left(Z^\alpha T_\alpha\right)(u_x),$$

for any $Z^\alpha \tilde{\partial}_\alpha + Y^\circ \dot{\tilde{\partial}}_\circ \in \Gamma\left((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E\right)$. Using this morphism, we obtain the tangent (ρ,η)-application $((\rho,\eta)\,T\pi, h \circ \pi)$ from $((\rho,\eta)\,TE, (\rho,\eta)\,\tau_E, E)$ to (F, ν, N).

Definition 5 *The kernel of the tangent (ρ,η)-application is denoted by*

$$\left(V\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_{E},E\right),$$

and it is called the vertical subbundle.

It is remarkable that $\left\{\dot{\tilde{\partial}}_{\circ}\right\}$ is a base of the $\mathcal{F}\left(E\right)$-module

$$\left(\Gamma\left(V\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_{E},E\right),+,\cdot\right).$$

The fiber bundles morphism (Π,π) defined by the commutative diagram

$$\begin{array}{ccc} V\left(\rho,\eta\right)TE & \overset{\Pi}{\longrightarrow} & E \\ \left(\rho,\eta\right)\tau_{E}\downarrow & & \downarrow\pi \\ E & \overset{\pi}{\longrightarrow} & M \end{array}$$

such that the components of the image of the vector $Y\dot{\tilde{\partial}}\left(u_{x}\right)$ are the real number $Y(u_{x})$ is called the *canonical projection fiber bundle morphism.*

Proposition 6 *The short sequence of vector bundles*

$$\begin{array}{ccccccccc} 0 & \hookrightarrow & V\left(\rho,\eta\right)TE & \hookrightarrow & \left(\rho,\eta\right)TE & \overset{(\rho,\eta)\pi!}{\longrightarrow} & \pi^{*}\left(h^{*}F\right) & \rightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ E & \overset{Id_{E}}{\longrightarrow} & E & \overset{Id_{E}}{\longrightarrow} & E & \overset{Id_{E}}{\longrightarrow} & E & \overset{Id_{E}}{\longrightarrow} & E \end{array}$$

is exact.

Definition 7 *A manifolds morphism $(\rho,\eta)\Gamma$ from $(\rho,\eta)TE$ to $V\left(\rho,\eta\right)TE$ defined by*

$$\left(\rho,\eta\right)\Gamma\left(Z^{\gamma}\tilde{\partial}_{\gamma}+Y^{\circ}\dot{\tilde{\partial}}_{\circ}\right)\left(u_{x}\right)=\left(Y^{\circ}+\left(\rho,\eta\right)\Gamma_{\gamma}^{\circ}Z^{\gamma}\right)\dot{\tilde{\partial}}_{\circ}\left(u_{x}\right),$$

such that the vector bundles morphism $((\rho,\eta)\Gamma,Id_{E})$ is a split to the left in the above exact sequence, is called (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M).

In particular case, the (ρ,Id_{M})-connection is called ρ-connection and it is denoted by $\rho\Gamma$ and in the classical case, the (Id_{TM},Id_{M})-connection is called connection and it is denoted by Γ.

Theorem 8 *If $(\rho,\eta)\Gamma$ is a (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M),* then its components satisfy the law of transformation

$$\left(\rho,\eta\right)\Gamma_{\gamma^{\prime}}^{\circ^{\prime}}=\frac{\partial y^{\circ\prime}}{\partial y^{\circ}}\left[\rho_{\gamma}^{k}\circ h\circ\pi\frac{\partial y^{\circ}}{\partial x^{k}}+\left(\rho,\eta\right)\Gamma_{\gamma}^{\circ}\right]\Lambda_{\gamma^{\prime}}^{\gamma}\circ h\circ\pi. \tag{4}$$

In the particular case of Lie algebroids, $(\eta, h) = (Id_M, Id_M)$, the above relation reduce to

$$\rho\Gamma^{\circ'}_{\gamma'} = \frac{\partial y^{\circ'}}{\partial y^{\circ}} \left[\rho^k_{\gamma} \circ \pi \frac{\partial y^{\circ}}{\partial x^k} + \rho\Gamma^{\circ}_{\gamma}\right] \Lambda^{\gamma}_{\gamma'} \circ \pi,$$

and in the classical case, $(\rho, \eta, h) = (Id_{TM}, Id_M, Id_M)$, the relations (4) reduce to

$$\Gamma^{\circ'}_{k'} = \frac{\partial y^{\circ'}}{\partial y^{\circ}} \left[\frac{\partial y^{\circ}}{\partial x^k} + \Gamma^{\circ}_{k}\right] \frac{\partial x^k}{\partial x^{k'}} \circ \tau_M.$$

Definition 9 *Let $(\rho, \eta)\Gamma$ be a (ρ, η)-connection for the Kaluza-Klein bundle (E, π, M). Then the kernel of the vector bundles morphism $((\rho, \eta)\Gamma, Id_E)$ is a vector subbundle of vector bundle $((\rho, \eta)TE, (\rho, \eta)\tau_E, E)$ which is called the horizontal subbundle and it is denoted by $(H(\rho, \eta)TE, (\rho, \eta)\tau_E, E)$.*

We put

$$\tilde{\delta}_{\alpha} = \tilde{\partial}_{\alpha} - (\rho, \eta)\Gamma_{\alpha}\dot{\tilde{\partial}}_{\circ} = T_{\alpha} \oplus \left(\left(\rho^i_{\alpha} \circ h \circ \pi\right)\partial_i - (\rho, \eta)\Gamma^{\circ}_{\alpha}\dot{\partial}_{\circ}\right).$$

It is easy to see that $\{\tilde{\delta}_{\alpha}\}$ is a basis for the $\mathcal{F}(E)$-module

$$\left(\Gamma\left(H(\rho, \eta)TE, (\rho, \eta)\tau_E, E\right), +, \cdot\right),$$

and $\left(\tilde{\delta}_{\alpha}, \dot{\tilde{\partial}}_{\circ}\right)$ is a basis for the $\mathcal{F}(E)$ -module

$$\left(\Gamma\left((\rho, \eta)TE, (\rho, \eta)\tau_E, E\right), +, \cdot\right),$$

which is called the *adapted (ρ, η)-base*. Moreover, the following equality holds

$$\Gamma(\tilde{\rho}, Id_E)\left(\tilde{\delta}_{\alpha}\right) = \left(\rho^i_{\alpha} \circ h \circ \pi\right)\partial_i - (\rho, \eta)\Gamma^{\circ}_{\alpha}\dot{\partial}_{\circ},$$

where $\left(\delta_i, \dot{\partial}_{\circ}\right)$ is the adapted base for the $\mathcal{F}(E)$-module $(\Gamma(TE, \tau_E, E), +, \cdot)$.

Theorem 10 *The following equalities hold*

$$\left[\tilde{\delta}_{\alpha}, \tilde{\delta}_{\beta}\right]_{(\rho,\eta)TE} = L^{\gamma}_{\alpha\beta} \circ (h \circ \pi)\tilde{\delta}_{\gamma} + (\rho, \eta, h)\mathbb{R}^{\circ}{}_{\alpha\beta}\dot{\tilde{\partial}}_{\circ},$$

$$\left[\tilde{\delta}_{\alpha}, \dot{\tilde{\partial}}_{\circ}\right]_{(\rho,\eta)TE} = \Gamma(\tilde{\rho}, Id_E)\left(\dot{\tilde{\partial}}_{\circ}\right)((\rho, \eta)\Gamma^{\circ}_{\alpha})\dot{\tilde{\partial}}_{\circ},$$

where

$$(\rho, \eta, h)\mathbb{R}^{\circ}{}_{\alpha\beta} = \Gamma(\tilde{\rho}, Id_E)\left(\tilde{\delta}_{\beta}\right)((\rho, \eta)\Gamma^{\circ}_{\alpha}) - \Gamma(\tilde{\rho}, Id_E)\left(\tilde{\delta}_{\alpha}\right)\left((\rho, \eta)\Gamma^{\circ}_{\beta}\right) + \left(L^{\gamma}_{\alpha\beta} \circ h \circ \pi\right)(\rho, \eta)\Gamma^{\circ}_{\gamma}.$$

If $(d\tilde{z}^{\alpha}, d\tilde{y}^{\circ})$ is the natural dual (ρ, η)-base of the natural (ρ, η)-base $\left(\tilde{\partial}_{\alpha}, \dot{\tilde{\partial}}_{\circ}\right)$, then it is easy to check that $(d\tilde{z}^{\alpha}, \delta\tilde{y}^{\circ})$ is the dual (ρ, η)-base of the adapted (ρ, η)-base $\left(\tilde{\delta}_{\alpha}, \dot{\tilde{\partial}}_{\circ}\right)$, where

$$\delta\tilde{y}^{\circ} = (\rho, \eta)\Gamma^{\circ}_{\alpha}d\tilde{z}^{\alpha} + d\tilde{y}^{\circ}.$$

The base $(d\tilde{z}^{\alpha}, \delta\tilde{y}^{\circ})$ is called the *adapted dual (ρ, η)-base.*

3.2 Distinguished linear (ρ,η)-connections

Let $(\rho,\eta)\,\Gamma$ be a (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M) and let

$$\left(\mathcal{T}\ {}^{p,1}_{q,1}\left((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E\right),+,\cdot\right),$$

be the $\mathcal{F}(E)$-module of $\binom{p,1}{q,1}$-tensor fields of the generalized tangent bundle

$$\left(H\left(\rho,\eta\right)TE,(\rho,\eta)\,\tau_E,E\right)\oplus\left(V\left(\rho,\eta\right)TE,(\rho,\eta)\,\tau_E,E\right).$$

An arbitrarily tensor field T with respect to the adapted (ρ,η)-base $\left(\tilde{\delta}_\alpha,\dot{\tilde{\partial}}_\circ\right)$ is written as

$$T=T^{\alpha_1...\alpha_p\circ}_{\beta_1...\beta_q\circ}\tilde{\delta}_{\alpha_1}\otimes...\otimes\tilde{\delta}_{\alpha_p}\otimes d\tilde{z}^{\beta_1}\otimes...\otimes d\tilde{z}^{\beta_q}\otimes\dot{\tilde{\partial}}_\circ\otimes\delta\tilde{y}^\circ.$$

Let $(\mathcal{T}\ ((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E),+,\cdot,\otimes)$ be the tensor fields algebra of generalized tangent bundle $((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E)$ and let

$$(X,T)\overset{(\rho,\eta)D}{\longrightarrow}(\rho,\eta)\,D_XT,$$

be a covariant (ρ,η)-derivative for the tensor algebra

$$(\mathcal{T}\ ((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E),+,\cdot,\otimes),$$

of the generalized tangent bundle $((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E)$ which preserves the horizontal and vertical interior differential systems by parallelism. The real local functions

$$\left((\rho,\eta)\,H^\alpha_{\beta\gamma},(\rho,\eta)\,H^\circ_{\circ\gamma},(\rho,\eta)\,V^\alpha_{\beta\circ},(\rho,\eta)\,V^\circ_{\circ\circ}\right),$$

defined by the following equalities:

$$\begin{aligned}&(\rho,\eta)\,D_{\tilde{\delta}_\gamma}\tilde{\delta}_\beta=(\rho,\eta)\,H^\alpha_{\beta\gamma}\tilde{\delta}_\alpha,\quad&&(\rho,\eta)\,D_{\tilde{\delta}_\gamma}\dot{\tilde{\partial}}_\circ=(\rho,\eta)\,H^\circ_{\circ\gamma}\dot{\tilde{\partial}}_\circ,\\&(\rho,\eta)\,D_{\dot{\tilde{\partial}}_\circ}\tilde{\delta}_\beta=(\rho,\eta)\,V^\alpha_{\beta\circ}\tilde{\delta}_\alpha,\quad&&(\rho,\eta)\,D_{\dot{\tilde{\partial}}_\circ}\dot{\tilde{\partial}}_\circ=(\rho,\eta)\,V^\circ_{\circ\circ}\dot{\tilde{\partial}}_\circ,\end{aligned}$$

are the components of a linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$ for the generalized tangent bundle $((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E)$ which is called the *distinguished linear (ρ,η)-connection.*

In the particular case of Lie algebroids, $h=Id_M=\eta$, we obtain the *distinguished linear ρ-connection.* The components of a distinguished linear ρ-connection $(\rho H,\rho V)$ are denoted by

$$\left(\rho H^\alpha_{\beta\gamma},\rho H^\circ_{\circ\gamma},\rho V^\alpha_{\beta\circ},\rho V^\circ_{\circ\circ}\right).$$

In addition, if $\rho=Id_{TM}$, then we obtain the classical *distinguished linear connection.* The components of a distinguished linear connection (H,V) are denoted by

$$\left(H^i_{jk},H^\circ_{\circ k},V^i_{j\circ},V^\circ_{\circ\circ}\right).$$

Theorem 11 *If* $((\rho,\eta)H,(\rho,\eta)V)$ is a distinguished linear (ρ,η)-connection for the generalized tangent bundle $((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E)$, then its components satisfy the change relations:

$$\begin{aligned}
(\rho,\eta)\,H_{\beta'\gamma'}^{\alpha'} &= \Lambda_{\alpha}^{\alpha'}\circ h\circ\pi\cdot\Big[\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)\left(\Lambda_{\beta}^{\alpha}\circ h\circ\pi\right)+\\
&\quad+(\rho,\eta)\,H_{\beta\gamma}^{\alpha}\cdot\Lambda_{\beta'}^{\beta}\circ h\circ\pi\Big]\cdot\Lambda_{\gamma'}^{\gamma}\circ h\circ\pi,\\
(\rho,\eta)\,H_{\circ'\gamma'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot\Big[\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)\left(\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\right)+\\
&\quad+(\rho,\eta)\,H_{\circ\gamma}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\Big]\cdot\Lambda_{\gamma'}^{\gamma}\circ h\circ\pi,\\
(\rho,\eta)\,V_{\beta'\circ'}^{\alpha'} &= \Lambda_{\alpha}^{\alpha'}\circ h\circ\pi\cdot(\rho,\eta)\,V_{\beta\circ}^{\alpha}\cdot\Lambda_{\beta'}^{\beta}\circ h\circ\pi\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}},\\
(\rho,\eta)\,V_{\circ'\circ'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot(\rho,\eta)\,V_{\circ\circ}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}.
\end{aligned}$$

In the particular case of Lie algebroids, $(\eta,h)=(Id_M,Id_M)$, we obtain

$$\begin{aligned}
\rho H_{\beta'\gamma'}^{\alpha'} &= \Lambda_{\alpha}^{\alpha'}\circ\pi\cdot\Big[\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)\left(\Lambda_{\beta'}^{\alpha}\circ\pi\right)+\rho H_{\beta\gamma}^{\alpha}\cdot\Lambda_{\beta'}^{\beta}\circ\pi\Big]\cdot\Lambda_{\gamma'}^{\gamma}\circ\pi,\\
\rho H_{\circ'\gamma'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot\Big[\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)\left(\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\right)+\rho H_{\circ\gamma}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\Big]\cdot\Lambda_{\gamma'}^{\gamma}\circ\pi,\\
\rho V_{\beta'\circ'}^{\alpha'} &= \Lambda_{\alpha'}^{\alpha}\circ\pi\cdot\rho V_{\beta\circ}^{\alpha}\cdot\Lambda_{\beta'}^{\beta}\circ\pi\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}},\\
\rho V_{\circ'\circ'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot\rho V_{\circ\circ}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}.
\end{aligned}$$

Also, in the classical case, $(\rho,\eta,h)=(Id_{TM},Id_M,Id_M)$, we deduce that the components of a distinguished linear connection (H,V) satisfy the change relations:

$$\begin{aligned}
H_{j'k'}^{i'} &= \tfrac{\partial x^{i\prime}}{\partial x^{i}}\circ\pi\cdot\Big[\tfrac{\delta}{\delta x^{k}}\left(\tfrac{\partial x^{i}}{\partial x^{j\prime}}\circ\pi\right)+H_{jk}^{i}\cdot\tfrac{\partial x^{j}}{\partial x^{j\prime}}\circ\pi\Big]\cdot\tfrac{\partial x^{k}}{\partial x^{k\prime}}\circ\pi,\\
H_{\circ'k'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot\Big[\tfrac{\delta}{\delta x^{k}}\left(\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\right)+H_{\circ k}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\Big]\cdot\tfrac{\partial x^{k}}{\partial x^{k\prime}}\circ\pi,\\
V_{j'\circ'}^{i'} &= \tfrac{\partial x^{i\prime}}{\partial x^{i}}\circ\pi\cdot V_{j\circ}^{i}\cdot\tfrac{\partial x^{j}}{\partial x^{j\prime}}\circ\pi\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}},\\
V_{\circ'\circ'}^{\circ'} &= \tfrac{\partial y^{\circ\prime}}{\partial y^{\circ}}\cdot V_{\circ\circ}^{\circ}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}\cdot\tfrac{\partial y^{\circ}}{\partial y^{\circ\prime}}.
\end{aligned}$$

Example 12 *The local real functions*

$$\left(\tfrac{\partial(\rho,\eta)\Gamma_{\gamma}^{\circ}}{\partial y^{\circ}},\tfrac{\partial(\rho,\eta)\Gamma_{\gamma}^{\circ}}{\partial y^{\circ}},0,0\right),$$

are the components of a distinguished linear (ρ,η)*-connection for the generalized tangent bundle* $((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E)$*, which is called the Berwald linear* (ρ,η)*-connection.*

Note that the Berwald linear (Id_{TM},Id_M)-connection is the usual *Berwald linear connection.*

Theorem 13 *If the generalized tangent bundle $((\rho,\eta)TE,(\rho,\eta)\tau_E,E)$ is endowed with a distinguished linear (ρ,η)-connection $((\rho,\eta)H,(\rho,\eta)V)$, then for any*

$$X=Z^{\alpha}\tilde{\delta}_{\alpha}+Y^{\circ}\dot{\tilde{\partial}}_{\circ}\in\Gamma((\rho,\eta)TE,(\rho,\eta)\tau_E,E),$$

and

$$T\in\mathcal{T}_{q1}^{p1}((\rho,\eta)TE,(\rho,\eta)\tau_E,E),$$

we obtain the formula:

$$\begin{aligned}(\rho,\eta)\,D_X&\left(T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\tilde{\delta}_{\alpha_1}\otimes\cdots\otimes\tilde{\delta}_{\alpha_p}\otimes d\tilde{z}^{\beta_1}\otimes\cdots\otimes d\tilde{z}^{\beta_q}\otimes\dot{\tilde{\partial}}_{\circ}\otimes\delta\tilde{y}^{\circ}\right)\\&=Z^{\gamma}T^{\alpha_1\ldots\alpha_p\circ}_{\beta_1\ldots\beta_q\circ|\gamma}\tilde{\delta}_{\alpha_1}\otimes\cdots\otimes\tilde{\delta}_{\alpha_p}\otimes d\tilde{z}^{\beta_1}\otimes\cdots\otimes d\tilde{z}^{\beta_q}\otimes\dot{\tilde{\partial}}_{\circ}\otimes\delta\tilde{y}^{\circ}\\&\quad+Y^{\circ}T^{\alpha_1\ldots\alpha_p\circ}_{\beta_1\ldots\beta_q\circ}\mid_{\circ}\tilde{\delta}_{\alpha_1}\otimes\cdots\otimes\tilde{\delta}_{\alpha_p}\otimes d\tilde{z}^{\beta_1}\otimes\cdots\otimes d\tilde{z}^{\beta_q}\otimes\dot{\tilde{\partial}}_{\circ}\otimes\delta\tilde{y}^{\circ},\end{aligned}$$

where

$$\begin{aligned}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ|\gamma}&=\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\ldots\beta_q\circ}\\&\quad+(\rho,\eta)\,H^{\alpha_1}_{\alpha\gamma}T^{\alpha\alpha_2\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}+\cdots+(\rho,\eta)\,H^{\alpha_p}_{\alpha\gamma}T^{\alpha_1\cdots\alpha_{p-1}\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad-(\rho,\eta)\,H^{\beta}_{\beta_1\gamma}T^{\alpha_1\cdots\alpha_p\circ}_{\beta\beta_2\cdots\beta_q\circ}-\cdots-(\rho,\eta)\,H^{\beta}_{\beta_q\gamma}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_{q-1}\beta\circ},\end{aligned}$$

and

$$\begin{aligned}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\mid_{\circ}&=\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad+(\rho,\eta)\,V^{\alpha_1}_{\alpha\circ}T^{\alpha\alpha_2\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}+\cdots+(\rho,\eta)\,V^{\alpha_p}_{\alpha\circ}T^{\alpha_1\cdots\alpha_{p-1}\alpha\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad-(\rho,\eta)\,V^{\beta}_{\beta_1\circ}T^{\alpha_1\cdots\alpha_p\circ}_{\beta\beta_2\cdots\beta_q\circ}-\cdots-(\rho,\eta)\,V^{\beta}_{\beta_q\circ}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_{q-1}\beta\circ}.\end{aligned}$$

In the particular case of Lie algebroids, $(\eta,h)=(Id_M,Id_M)$, we obtain

$$\begin{aligned}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ|\gamma}&=\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\ldots\beta_q\circ}\\&\quad+\rho H^{\alpha_1}_{\alpha\gamma}T^{\alpha\alpha_2\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}+\cdots+\rho H^{\alpha_p}_{\alpha\gamma}T^{\alpha_1\cdots\alpha_{p-1}\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad-\rho H^{\beta}_{\beta_1\gamma}T^{\alpha_1\cdots\alpha_p\circ}_{\beta\beta_2\cdots\beta_q\circ}-\cdots-\rho H^{\beta}_{\beta_q\gamma}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_{q-1}\beta\circ},\end{aligned}$$

and

$$\begin{aligned}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\mid_{\circ}&=\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad+\rho V^{\alpha_1}_{\alpha\circ}T^{\alpha\alpha_2\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}+\cdots+\rho V^{\alpha_p}_{\alpha\circ}T^{\alpha_1\cdots\alpha_{p-1}\alpha\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad-\rho V^{\beta}_{\beta_1\circ}T^{\alpha_1\cdots\alpha_p\circ}_{\beta\beta_2\cdots\beta_q\circ}-\cdots-\rho V^{\beta}_{\beta_q\circ}T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_{q-1}\beta\circ}.\end{aligned}$$

Moreover, in the classical case, $(\rho,\eta,h)=(Id_{TM},Id_M,Id_M)$, we obtain

$$\begin{aligned}T^{i_1\cdots i_p\circ}_{j_1\cdots j_q\circ|k}&=\delta_k\left(T^{i_1\cdots i_p\circ}_{j_1\cdots j_q\circ}\right)\\&\quad+H^{i_1}_{ik}T^{ii_2\cdots i_p\circ}_{j_1\cdots j_q\circ}+\cdots+H^{i_p}_{ik}T^{i_1\cdots i_{p-1}i\circ}_{\beta_1\cdots\beta_q\circ}\\&\quad-H^{j}_{j_1k}T^{i_1\cdots i_p\circ}_{jj_2\cdots j_q\circ}-\cdots-H^{j}_{j_qk}T^{\alpha_1\cdots\alpha_p\circ}_{j_1\cdots j_{q-1}j\circ},\end{aligned}$$

and

$$T^{i_1\cdots i_p\circ}_{j_1\cdots j_q\circ}\mid_\circ = \dot{\partial}_\circ\left(T^{\alpha_1\cdots\alpha_p\circ}_{\beta_1\cdots\beta_q\circ}\right)$$
$$+V^{i_1}_{i\circ}T^{ii_2\cdots i_p\circ}_{j_1\cdots j_q\circ}+\cdots+V^{i_p}_{i\circ}T^{i_1\cdots i_{p-1}i\circ}_{\beta_1\cdots\beta_q\circ}$$
$$-V^{j}_{j_1\circ}T^{i_1\cdots i_p\circ}_{jj_2\cdots j_q\circ}-\cdots-V^{j}_{j_q\circ}T^{i_1\cdots i_p\circ}_{j_1\cdots j_{q-1}j\circ}.$$

4 The (g,h)-lift of a differentiable curve

In this section we use the Kaluza-Klein bundle such that the fibre K is an 1-dimensional real vector space. Thus we obtain the Kaluza-Klein vector bundle (E,π,M).

Let $c:I\to M$ be a differentiable curve. Then

$$\left(E_{|\operatorname{Im}(\eta\circ h\circ c)},\pi_{|\operatorname{Im}(\eta\circ h\circ c)},\operatorname{Im}\left(\eta\circ h\circ c\right)\right),$$

is a vector subbundle of (E,π,M).

Definition 14 *Let*

$$\begin{array}{ccc} I & \xrightarrow{\dot{c}} & E_{|\operatorname{Im}(\eta\circ h\circ c)}, \\ t & \longmapsto & y^\circ(t)\,s_\circ\left(\eta\circ h\circ c\left(t\right)\right), \end{array}$$

be a differentiable curve. If there exists a manifolds morphism $g:E\to F$ such that the following conditions are hold:

1) (g,h) is a vector bundle morphism from (E,π,M) to (F,ν,N),

2) $\rho\circ g\circ\dot{c}\left(t\right)=\dfrac{d\left(\eta\circ h\circ c\right)^i(t)}{dt}\dfrac{\partial}{\partial x^i}\left(\left(\eta\circ h\circ c\right)(t)\right)$, for any $t\in I$,

then $\dot{c}$ is called the (g,h)-lift of the differentiable curve c.

Remark 15 *The second condition is equivalent with the following:*

$$\rho^i_\alpha\left(\eta\circ h\circ c\left(t\right)\right)\cdot g^\alpha_\circ\left(h\circ c\left(t\right)\right)\cdot y^\circ\left(t\right)=\tfrac{d(\eta\circ h\circ c)^i(t)}{dt},\ i\in 1,\ldots,4.$$

Definition 16 *If $\dot{c}:I\longrightarrow E_{|\operatorname{Im}(\eta\circ h\circ c)}$, is a differentiable (g,h)-lift of the differentiable curve c, then the section*

$$\begin{array}{ccc} \operatorname{Im}\left(\eta\circ h\circ c\right) & \xrightarrow{u(c,\dot{c})} & E_{|\operatorname{Im}(\eta\circ h\circ c)}, \\ \eta\circ h\circ c\left(t\right) & \longmapsto & \dot{c}\left(t\right), \end{array}$$

is called the canonical section associated to $(c,\dot{c})$.

Definition 17 *If (g,h) has the components*

$$g^\alpha_\circ,\quad \alpha\in 1,\ldots,p,$$

such that for any local vector chart (V,t_V) of (F,ν,N), there exists the real functions

$$V\xrightarrow{\tilde{g}^\circ_\alpha}\mathbb{R},\quad \alpha\in 1,\ldots,p,$$

such that

$$\tilde{g}_{\beta}^{\circ}(\varkappa)\cdot g_{\circ}^{\alpha}(\varkappa)=\delta_{\beta}^{\alpha},$$

for any $\varkappa\in V$*, then we say that the* (g,h) *is locally invertible.*

Definition 18 *If* $\dot{c}:I\longrightarrow E_{|\operatorname{Im}(\eta\circ h\circ c)}$ *is a differentiable* (g,h)*-lift of differentiable curve* c*, such that its component function* (y°) *is solution for the differentiable system of equations:*

$$\tfrac{du^{\circ}}{dt}+(\rho,\eta)\,\Gamma_{\alpha}^{\circ}\circ u(c,\dot{c})\circ(\eta\circ h\circ c)\cdot g_{\circ}^{\alpha}\circ h\circ c\cdot u^{\circ}=0,$$

then we say that the (g,h)*-lift* $\dot{c}$ *is parallel with respect to the* (ρ,η)*-connection* $(\rho,\eta)\,\Gamma$*.*

4.1 The lift of accelerations for a differentiable curve

Let

$$\begin{array}{ccc} I & \overset{\dot{c}}{\longrightarrow} & E_{|\operatorname{Im}(\eta\circ h\circ c)},\\ t & \longmapsto & y^{\circ}(t)\,s_{\circ}(\eta\circ h\circ c(t)),\end{array}$$

be the (g,h)-lift of differentiable curve $I\overset{c}{\longrightarrow}M$.

Definition 19 *The differentiable curve* $\ddot{c}:I\longrightarrow(\rho,\eta)\,TE_{|\operatorname{Im}\dot{c}}$ *defined by*

$$\ddot{c}(t)=(g_{\circ}^{\alpha}\circ h\circ c(t)\cdot y^{\circ}(t))\,\tilde{\partial}_{\alpha}(\dot{c}(t))+\tfrac{dy^{\circ}(t)}{dt}\dot{\tilde{\partial}}_{\circ}(\dot{c}(t)),$$

is called the differentiable (g,h)*-lift of accelerations of the differentiable curve* c*. Moreover, the section*

$$\begin{array}{ccc}\operatorname{Im}(\dot{c}) & \overset{u(c,\dot{c},\ddot{c})}{\longrightarrow} & (\rho,\eta)\,TE_{|\operatorname{Im}(\dot{c})},\\ \dot{c}(t) & \longmapsto & (g_{\circ}^{\alpha}\circ h\circ c(t)\cdot y^{\circ}(t))\,\tilde{\partial}_{\alpha}(\dot{c}(t))+\dfrac{dy^{\circ}(t)}{dt}\dot{\tilde{\partial}}_{\circ}(\dot{c}(t)),\end{array}$$

is called the canonical section associated to the triple $(c,\dot{c},\ddot{c})$*.*

In the adapted (ρ,η)-base $\left(\tilde{\delta}_{\alpha},\dot{\tilde{\partial}}_{\circ}\right)$, we can rewrite the above equation as follows:

$$\begin{aligned}u(c,\dot{c},\ddot{c})(\dot{c}(t)) &= (g_{\circ}^{\alpha}\circ h\circ c(t)\,y^{\circ}(t))\,\tilde{\delta}_{\alpha}(\dot{c}(t))+\frac{dy^{\circ}(t)}{dt}\dot{\tilde{\partial}}_{\circ}(\dot{c}(t))\\ &\quad+(\rho,\eta)\,\Gamma_{\alpha}^{\circ}\circ u(c,\dot{c})\circ\eta\circ h\circ c(t)\cdot(g_{\circ}^{\alpha}\circ h\circ c(t)\,y^{\circ}(t))\,\dot{\tilde{\partial}}_{\circ}(\dot{c}(t)),\end{aligned}$$

where $t\in I$. It is easy to check that $u(c,\dot{c},\ddot{c})(\dot{c}(t))\in H(\rho,\eta)\,TE_{|\operatorname{Im}(\dot{c})}$ if and only if the component function y° is solution for the differentiable equations

$$\tfrac{du^{\circ}}{dt}+(\rho,\eta)\,\Gamma_{\alpha}^{\circ}\circ u(c,\dot{c})\circ\eta\circ h\circ c\cdot(g_{\circ}^{\alpha}\circ h\circ c)\cdot u^{\circ}=0.$$

Definition 20 *If the component functions* $((g_{\circ}^{\alpha} \circ h \circ c)\, y^{\circ})$ *are solutions for the differentiable system of equations*

$$\tfrac{dz^{\alpha}}{dt} + (\rho,\eta)\, H_{\beta\gamma}^{\alpha} \circ u\,(c,\dot{c}) \circ \eta \circ h \circ c \cdot z^{\beta} \cdot z^{\gamma} = 0,\ \alpha \in 1, \cdots, p,$$

then the differentiable curve $\dot{c}$ *is called horizontal parallel with respect to the distinguished linear* (ρ,η)*-connection* $((\rho,\eta)\, H, (\rho,\eta)\, V)$*. Also, if the component function* y° *is a solution for the differentiable system of equations*

$$\tfrac{du^{\circ}}{dt} + (\rho,\eta)\, V_{\circ\circ}^{\circ} \circ u\,(c,\dot{c}) \circ \eta \circ h \circ c \cdot u^{\circ} \cdot u^{\circ} = 0,$$

then the differentiable curve $\dot{c}$ *is called vertical parallel with respect to the distinguished linear* (ρ,η)*-connection* $((\rho,\eta)\, H, (\rho,\eta)\, V)$*.*

5 The (ρ,η,h)-torsion and the (ρ,η,h)-curvature of a distinguished linear (ρ,η)-connection

Let $(\rho,\eta)\Gamma$ be a (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M) and let $((\rho,\eta)H,(\rho,\eta)V)$ be a distinguished linear (ρ,η)-connection for the Lie algebroid generalized tangent bundle

$$\left(((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E), [,]_{(\rho,\eta)TE}, (\tilde{\rho}, Id_E)\right).$$

Definition 21 *The application*

$$\begin{array}{ccc} \Gamma\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right)^2 & \overset{(\rho,\eta,h)\mathbb{T}}{\longrightarrow} & \Gamma\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right), \\ (X,Y) & \longmapsto & (\rho,\eta)\, \mathbb{T}\,(X,Y), \end{array}$$

defined by

$$(\rho,\eta,h)\, \mathbb{T}\,(X,Y) = (\rho,\eta)\, D_X Y - (\rho,\eta)\, D_Y X - [X,Y]_{(\rho,\eta)TE},$$

for any $X, Y \in \Gamma\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right)$*, is called the* (ρ,η,h)*-torsion associated to distinguished linear* (ρ,η)*-connection* $((\rho,\eta)\, H, (\rho,\eta)\, V)$*. Moreover, the applications*

$$\mathcal{H}\,(\rho,\eta,h)\, \mathbb{T}\,(\mathcal{H}\,(\cdot), \mathcal{H}\,(\cdot)),\ \mathcal{V}\,(\rho,\eta,h)\, \mathbb{T}\,(\mathcal{H}\,(\cdot), \mathcal{H}\,(\cdot)), \ldots, \mathcal{V}\,(\rho,\eta,h)\, \mathbb{T}\,(\mathcal{V}\,(\cdot), \mathcal{V}\,(\cdot))$$

are called $\mathcal{H}\,(\mathcal{H}\mathcal{H}),\ \mathcal{V}\,(\mathcal{H}\mathcal{H}), \ldots, \mathcal{V}\,(\mathcal{V}\mathcal{V})$ (ρ,η,h)*-torsions associated to distinguished linear* (ρ,η)*-connection* $((\rho,\eta)\, H, (\rho,\eta)\, V)$*.*

Proposition 22 *The* (ρ,η,h)*-torsion* $(\rho,\eta,h)\, \mathbb{T}$ *associated to distinguished linear* (ρ,η)*-connection* $((\rho,\eta)\, H, (\rho,\eta)\, V)$*, is* $\mathbb{R}$*-bilinear and antisymmetric in the lower indices.*

Using the notations

$$\begin{aligned}
\mathcal{H}(\rho,\eta,h)\,\mathbb{T}\left(\tilde{\delta}_\gamma,\tilde{\delta}_\beta\right) &= (\rho,\eta,h)\,\mathbb{T}^{\alpha}{}_{\beta\gamma}\tilde{\delta}_\alpha,\\
\mathcal{V}(\rho,\eta,h)\,\mathbb{T}\left(\tilde{\delta}_\gamma,\tilde{\delta}_\beta\right) &= (\rho,\eta,h)\,\mathbb{T}^{\circ}{}_{\beta\gamma}\dot{\tilde{\partial}}_\circ,\\
\mathcal{H}(\rho,\eta,h)\,\mathbb{T}\left(\dot{\tilde{\partial}}_\circ,\tilde{\delta}_\beta\right) &= (\rho,\eta,h)\,\mathbb{P}^{\alpha}{}_{\beta\circ}\tilde{\delta}_\alpha,\\
\mathcal{V}(\rho,\eta,h)\,\mathbb{T}\left(\dot{\tilde{\partial}}_\circ,\tilde{\delta}_\beta\right) &= (\rho,\eta,h)\,\mathbb{P}^{\circ}{}_{\beta\circ}\dot{\tilde{\partial}}_\circ,\\
\mathcal{V}(\rho,\eta,h)\,\mathbb{T}\left(\dot{\tilde{\partial}}_\circ,\dot{\tilde{\partial}}_\circ\right) &= (\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\circ}\dot{\tilde{\partial}}_\circ,
\end{aligned}$$

we can prove the following

Theorem 23 *The (ρ,η,h)-torsion $(\rho,\eta,h)\,\mathbb{T}$ associated to the distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$, is characterized by the tensor fields with local components:*

$$\begin{aligned}
(\rho,\eta,h)\,\mathbb{T}^{\alpha}{}_{\beta\gamma} &= (\rho,\eta)\,H^{\alpha}_{\beta\gamma} - (\rho,\eta)\,H^{\alpha}_{\gamma\beta} - L^{\alpha}_{\beta\gamma}\circ h\circ\pi,\\
(\rho,\eta,h)\,\mathbb{T}^{\circ}{}_{\beta\gamma} &= (\rho,\eta,h)\,\mathbb{R}^{\circ}{}_{\beta\gamma},\\
(\rho,\eta,h)\,\mathbb{P}^{\alpha}{}_{\beta\circ} &= (\rho,\eta)\,V^{\alpha}_{\beta\circ},\\
(\rho,\eta,h)\,\mathbb{P}^{\circ}{}_{\beta\circ} &= \frac{\partial}{\partial y^\circ}\left((\rho,\eta)\,\Gamma^{\circ}_{\beta}\right) - (\rho,\eta)\,H^{\circ}_{\circ\beta},\\
(\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\circ} &= (\rho,\eta)\,V^{\circ}_{\circ\circ} - (\rho,\eta)\,V^{\circ}_{\circ\circ} = 0.
\end{aligned}$$

In particular, when $(\rho,\eta,h) = (Id_{TM}, Id_M, Id_M)$, we deduce the following local components of torsion associated to distinguished linear connection (H,V):

$$\begin{aligned}
\mathbb{T}^{i}{}_{jk} &= H^{i}_{jk} - H^{i}_{kj}, & \mathbb{T}^{\circ}{}_{jk} &= \mathbb{R}^{\circ}{}_{jk},\\
\mathbb{P}^{i}{}_{j\circ} &= V^{i}_{j\circ}, & \mathbb{P}^{\circ}{}_{j\circ} &= \frac{\partial\Gamma^{\circ}_{j}}{\partial y^\circ} - H^{\circ}_{\circ j},\\
\mathbb{S}^{\circ}{}_{\circ\circ} &= V^{\circ}_{\circ\circ} - V^{\circ}_{\circ\circ} = 0.
\end{aligned}$$

Definition 24 *The application*

$$\begin{array}{ccc}
\left(\Gamma\left((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E\right)\right)^3 & \xrightarrow{(\rho,\eta,h)\mathbb{R}} & \Gamma\left((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E\right),\\
((Y,Z),X) & \longmapsto & (\rho,\eta,h)\,\mathbb{R}\left((Y,Z),X\right),
\end{array}$$

defined by

$$\begin{aligned}
(\rho,\eta,h)\,\mathbb{R}(Y,Z)\,X = (\rho,\eta)\,D_Y\left((\rho,\eta)\,D_Z X\right) - (\rho,\eta)\,D_Z\left((\rho,\eta)\,D_Y X\right)\\
- (\rho,\eta)\,D_{[Y,Z]_{(\rho,\eta)TE}}X,
\end{aligned}$$

for any $X,Y,Z\in\Gamma\left((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E\right)$, is called the (ρ,η,h)-curvature associated to distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$.

Proposition 25 *The (ρ,η,h)-curvature $(\rho,\eta,h)\,\mathbb{R}$ associated to distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$, is $\mathbb{R}$-linear in each argument and antisymmetric in the first two arguments.*

Using the notations

$$\begin{aligned}
(\rho,\eta,h)\,\mathbb{R}\left(\tilde{\delta}_{\varepsilon},\tilde{\delta}_{\gamma}\right)\tilde{\delta}_{\beta} &= (\rho,\eta,h)\,\mathbb{R}^{\alpha}{}_{\beta\ \gamma\varepsilon}\tilde{\delta}_{\alpha},\\
(\rho,\eta,h)\,\mathbb{R}\left(\tilde{\delta}_{\varepsilon},\tilde{\delta}_{\gamma}\right)\dot{\tilde{\partial}}_{\circ} &= (\rho,\eta,h)\,\mathbb{R}^{\circ}{}_{\circ\ \gamma\varepsilon}\dot{\tilde{\partial}}_{\circ},\\
(\rho,\eta,h)\,\mathbb{R}\left(\dot{\tilde{\partial}}_{\circ},\tilde{\delta}_{\gamma}\right)\tilde{\delta}_{\varepsilon} &= (\rho,\eta,h)\,\mathbb{P}^{\alpha}{}_{\varepsilon\ \gamma\circ}\tilde{\delta}_{\alpha},\\
(\rho,\eta,h)\,\mathbb{R}\left(\dot{\tilde{\partial}}_{\circ},\tilde{\delta}_{\gamma}\right)\dot{\tilde{\partial}}_{\circ} &= (\rho,\eta,h)\,\mathbb{P}^{\circ}{}_{\circ\ \gamma\circ}\dot{\tilde{\partial}}_{\circ},\\
(\rho,\eta,h)\,\mathbb{R}\left(\dot{\tilde{\partial}}_{\circ},\dot{\tilde{\partial}}_{\circ}\right)\tilde{\delta}_{\beta} &= (\rho,\eta,h)\,\mathbb{S}^{\alpha}{}_{\beta\ \circ\circ}\tilde{\delta}_{\alpha},\\
(\rho,\eta,h)\,\mathbb{R}\left(\dot{\tilde{\partial}}_{\circ},\dot{\tilde{\partial}}_{\circ}\right)\dot{\tilde{\partial}}_{\circ} &= (\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\ \circ\circ}\dot{\tilde{\partial}}_{\circ},
\end{aligned}$$

we derive the following

Theorem 26 *The (ρ,η,h)-curvature $(\rho,\eta,h)\,\mathbb{R}$ associated to distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$, is characterized by the tensor fields with the local components:*

$$\left\{\begin{array}{l}
(\rho,\eta,h)\mathbb{R}^{\alpha}{}_{\beta\ \gamma\varepsilon}=\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\varepsilon}\right)(\rho,\eta)H^{\alpha}_{\beta\gamma}-\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\gamma}\right)(\rho,\eta)H^{\alpha}_{\beta\varepsilon}\\
\qquad\qquad +(\rho,\eta)H^{\alpha}_{\theta\varepsilon}(\rho,\eta)H^{\theta}_{\beta\gamma}-(\rho,\eta)H^{\alpha}_{\theta\gamma}(\rho,\eta)H^{\theta}_{\beta\varepsilon}\\
\qquad\qquad -(\rho,\eta,h)\mathbb{R}^{\circ}{}_{\gamma\varepsilon}(\rho,\eta)H^{\alpha}_{\beta\circ}-L^{\theta}_{\gamma\varepsilon}\circ h\circ\pi(\rho,\eta)H^{\alpha}_{\beta\theta},\\
(\rho,\eta,h)\mathbb{R}^{\circ}{}_{\circ\ \gamma\varepsilon}=\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\varepsilon}\right)(\rho,\eta)H^{\circ}_{\circ\gamma}-\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\gamma}\right)(\rho,\eta)H^{\circ}_{\circ\varepsilon}\\
\qquad\qquad +(\rho,\eta)H^{\circ}_{\circ\varepsilon}(\rho,\eta)H^{\circ}_{\circ\gamma}-(\rho,\eta)H^{\circ}_{\circ\gamma}(\rho,\eta)H^{\circ}_{\circ\varepsilon}\\
\qquad\qquad -(\rho,\eta,h)\mathbb{R}^{\circ}{}_{\varepsilon\gamma}(\rho,\eta)V^{\circ}_{\circ\circ}-L^{\theta}_{\gamma\varepsilon}\circ h\circ\pi(\rho,\eta)V^{\circ}_{\circ\theta},
\end{array}\right.$$

$$\left\{\begin{array}{l}
(\rho,\eta,h)\mathbb{P}^{\alpha}{}_{\varepsilon\ \gamma\circ}=\Gamma(\tilde{\rho},Id_E)(\dot{\tilde{\partial}}_{\circ})(\rho,\eta)H^{\alpha}_{\varepsilon\gamma}-\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\gamma}\right)(\rho,\eta)V^{\alpha}_{\varepsilon\circ}\\
\qquad\qquad +(\rho,\eta)V^{\alpha}_{\theta\circ}(\rho,\eta)H^{\theta}_{\varepsilon\gamma}-(\rho,\eta)H^{\alpha}_{\theta\gamma}(\rho,\eta)V^{\theta}_{\varepsilon\circ}\\
\qquad\qquad +\dot{\partial}_{\circ}\left((\rho,\eta)\,\Gamma^{\circ}_{\gamma}\right)(\rho,\eta)\,V^{\alpha}_{\varepsilon\circ},\\
(\rho,\eta,h)\mathbb{P}^{\circ}{}_{\circ\ \gamma\circ}=\Gamma(\tilde{\rho},Id_E)\left(\dot{\tilde{\partial}}_{\circ}\right)(\rho,\eta)H^{\circ}_{\circ\gamma}-\\
\qquad\qquad -\Gamma(\tilde{\rho},Id_E)\left(\tilde{\delta}_{\gamma}\right)(\rho,\eta)V^{\circ}_{\circ\circ}+(\rho,\eta)V^{\circ}_{\circ\circ}(\rho,\eta)H^{\circ}_{\circ\gamma}-\\
\qquad\qquad -(\rho,\eta)H^{\circ}_{\circ\gamma}(\rho,\eta)V^{\circ}_{\circ\circ}+\dot{\partial}_{\circ}(\left(\rho,\eta\right)\Gamma^{\circ}_{\gamma})\,(\rho,\eta)V^{\circ}_{\circ\circ},
\end{array}\right.$$

$$\begin{cases} (\rho,\eta,h)\,\mathbb{S}^{\alpha}_{\ \beta\ \circ\circ} = \Gamma\left(\tilde{\rho}, Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)(\rho,\eta)\,V^{\alpha}_{\beta\circ} \\ \qquad\qquad -\Gamma\left(\tilde{\rho}, Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)(\rho,\eta)\,V^{\alpha}_{\beta\circ} + (\rho,\eta)\,V^{\alpha}_{\theta\circ}\,(\rho,\eta)\,V^{\theta}_{\beta\circ} \\ \qquad\qquad -(\rho,\eta)\,V^{\alpha}_{\theta\circ}\,(\rho,\eta)\,V^{\theta}_{\beta\circ} = 0, \\ (\rho,\eta,h)\,\mathbb{S}^{\circ}_{\ \circ\ \circ\circ} = \Gamma\left(\tilde{\rho}, Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)(\rho,\eta)\,V^{\circ}_{\circ\circ} \\ \qquad\qquad -\Gamma\left(\tilde{\rho}, Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)(\rho,\eta)\,V^{\circ}_{\circ\circ} + (\rho,\eta)\,V^{\circ}_{\circ\circ}\,(\rho,\eta)\,V^{\circ}_{\circ\circ} \\ \qquad\qquad -(\rho,\eta)\,V^{\circ}_{\circ\circ}\,(\rho,\eta)\,V^{\circ}_{\circ\circ} = 0. \end{cases}$$

In particular, when $(\rho,\eta,h) = (Id_{TM}, Id_M, Id_M)$, we obtain the following local components of the curvature associated to distinguished linear connection (H,V):

$$\begin{aligned} \mathbb{R}^{i}_{\ j\ kl} &= \delta_l\left(H^i_{jk}\right) - \delta_k\left(H^i_{jl}\right) + H^i_{hl}H^h_{jk} - H^i_{hk}H^h_{jl} - \mathbb{R}^{\circ}_{\ kl}H^i_{j\circ}, \\ \mathbb{R}^{\circ}_{\ \circ\ kl} &= \delta_l\left(H^{\circ}_{\circ k}\right) - \delta_k\left(H^{\circ}_{\circ l}\right) + H^{\circ}_{\circ l}H^{\circ}_{\circ k} - H^{\circ}_{\circ k}H^{\circ}_{\circ l} - \mathbb{R}^{\circ}_{\ lk}V^{\circ}_{\circ\circ}, \\ \mathbb{P}^{i}_{\ l\ k\circ} &= \dot{\partial}_{\circ}\left(H^i_{lk}\right) - \delta_k\left(V^i_{l\circ}\right) + V^i_{h\circ}H^h_{lk} - H^i_{hk}V^h_{l\circ} + \dot{\partial}_{\circ}\left(\Gamma^{\circ}_{k}\right)V^i_{l\circ}, \\ \mathbb{P}^{\circ}_{\ \circ\ k\circ} &= \dot{\partial}_{\circ}\left(H^{\circ}_{\circ k}\right) - \delta_k\left(V^{\circ}_{\circ\circ}\right) + V^{\circ}_{\circ\circ}H^{\circ}_{\circ k} - H^{\circ}_{\circ k}V^{\circ}_{\circ\circ} + \dot{\partial}_{\circ}\left(\Gamma^{\circ}_{k}\right)V^{\circ}_{\circ\circ}, \\ \mathbb{S}^{i}_{\ j\ \circ\circ} &= \dot{\partial}_{\circ}\left(V^i_{j\circ}\right) - \dot{\partial}_{\circ}\left(V^i_{j\circ}\right) + V^i_{h\circ}V^h_{j\circ} - V^i_{h\circ}V^h_{j\circ} = 0, \\ \mathbb{S}^{\circ}_{\ \circ\ \circ\circ} &= \dot{\partial}_{\circ}\left(V^{\circ}_{\circ\circ}\right) - \dot{\partial}_{\circ}\left(V^{\circ}_{\circ\circ}\right) + V^{\circ}_{\circ\circ}V^{\circ}_{\circ\circ} - V^{\circ}_{\circ\circ}V^{\circ}_{\circ\circ} = 0. \end{aligned}$$

Definition 27 *The tensor field*

$$\begin{aligned} \mathbf{Ric}\left((\rho,\eta)\,H,(\rho,\eta)\,V\right) &= (\rho,\eta,h)\,\mathbb{R}_{\ \alpha\ \beta}d\tilde{z}^{\alpha}\otimes d\tilde{z}^{\beta} + (\rho,\eta,h)\,\mathbb{P}_{\ \alpha\ \circ}d\tilde{z}^{\alpha}\otimes\delta\tilde{y}^{\circ} \\ &\quad + (\rho,\eta,h)\,\mathbb{P}_{\ \circ\ \beta}\delta\tilde{y}^{\circ}\otimes d\tilde{z}^{\beta} + (\rho,\eta,h)\,\mathbb{S}_{\ \circ\ \circ}\delta\tilde{y}^{\circ}\otimes\delta\tilde{y}^{\circ}, \end{aligned}$$

is called the Ricci tensor field associated to distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$, where

$$\begin{aligned} (\rho,\eta,h)\,\mathbb{R}_{\ \alpha\ \beta} &= (\rho,\eta,h)\,\mathbb{R}^{\gamma}_{\ \alpha\ \beta\gamma}, \quad (\rho,\eta,h)\,\mathbb{P}_{\ \alpha\ \circ} = (\rho,\eta,h)\,\mathbb{P}^{\beta}_{\ \alpha\ \beta\circ}, \\ (\rho,\eta,h)\,\mathbb{P}_{\ \circ\ \beta} &= (\rho,\eta,h)\,\mathbb{P}^{\circ}_{\ \circ\ \beta\circ}, \quad (\rho,\eta,h)\,\mathbb{S}_{\ \circ\ \circ} = (\rho,\eta,h)\,\mathbb{S}^{\circ}_{\ \circ\ \circ\circ}. \end{aligned}$$

5.1 Identities of Cartan and Bianchi type

Let $(\rho,\eta)\Gamma$ be a (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M) and let $((\rho,\eta)H,(\rho,\eta)V)$ be a distinguished linear (ρ,η)-connection for the Lie algebroid generalized tangent bundle

$$\left(\left((\rho,\eta)\,TE,(\rho,\eta)\,\tau_E,E\right),[,]_{(\rho,\eta)TE},(\tilde{\rho},Id_E)\right).$$

Using the definition of (ρ,η,h)-curvature associated to the distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$, we have the following

Theorem 28 *The following equalities hold*

$$\begin{cases} (\rho,\eta)\, D_{\mathcal{H}X}\,(\rho,\eta)\, D_{\mathcal{H}Y}\mathcal{H}Z-(\rho,\eta)\, D_{\mathcal{H}Y}\,(\rho,\eta)\, D_{\mathcal{H}X}\mathcal{H}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{H}X,\mathcal{H}Y)\,\mathcal{H}Z+(\rho,\eta)\, D_{\mathcal{H}[\mathcal{H}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{H}Z \\ +(\rho,\eta)\, D_{\mathcal{V}[\mathcal{H}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{H}Z, \\ (\rho,\eta)\, D_{\mathcal{V}X}\,(\rho,\eta)\, D_{\mathcal{H}Y}\mathcal{H}Z-(\rho,\eta)\, D_{\mathcal{H}Y}\,(\rho,\eta)\, D_{\mathcal{V}X}\mathcal{H}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{V}X,\mathcal{H}Y)\,\mathcal{H}Z+(\rho,\eta)\, D_{\mathcal{H}[\mathcal{V}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{H}Z \\ +(\rho,\eta)\, D_{\mathcal{V}[\mathcal{V}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{H}Z, \\ (\rho,\eta)\, D_{\mathcal{V}X}\,(\rho,\eta)\, D_{\mathcal{V}Y}\mathcal{H}Z-(\rho,\eta)\, D_{\mathcal{V}Y}\,(\rho,\eta)\, D_{\mathcal{V}X}\mathcal{H}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{V}X,\mathcal{V}Y)\,\mathcal{H}Z+(\rho,\eta)\, D_{\mathcal{V}[\mathcal{V}X,\mathcal{V}Y]_{(\rho,\eta)TE}}\mathcal{H}Z, \end{cases} \tag{5}$$

and

$$\begin{cases} (\rho,\eta)\, D_{\mathcal{H}X}\,(\rho,\eta)\, D_{\mathcal{H}Y}\mathcal{V}Z-(\rho,\eta)\, D_{\mathcal{H}Y}\,(\rho,\eta)\, D_{\mathcal{H}X}\mathcal{V}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{H}X,\mathcal{H}Y)\,\mathcal{V}Z+(\rho,\eta)\, D_{\mathcal{H}[\mathcal{H}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{V}Z \\ +(\rho,\eta)\, D_{\mathcal{V}[\mathcal{H}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{V}Z, \\ (\rho,\eta)\, D_{\mathcal{V}X}\,(\rho,\eta)\, D_{\mathcal{H}Y}\mathcal{V}Z-(\rho,\eta)\, D_{\mathcal{H}Y}\,(\rho,\eta)\, D_{\mathcal{V}X}\mathcal{V}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{V}X,\mathcal{H}Y)\,\mathcal{V}Z+(\rho,\eta)\, D_{h[\mathcal{V}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{V}Z \\ +(\rho,\eta)\, D_{\mathcal{V}[\mathcal{V}X,\mathcal{H}Y]_{(\rho,\eta)TE}}\mathcal{V}Z, \\ (\rho,\eta)\, D_{\mathcal{V}X}\,(\rho,\eta)\, D_{\mathcal{V}Y}\mathcal{V}Z-(\rho,\eta)\, D_{\mathcal{V}Y}\,(\rho,\eta)\, D_{\mathcal{V}X}\mathcal{V}Z \\ =(\rho,\eta,h)\,\mathbb{R}\,(\mathcal{V}X,\mathcal{V}Y)\,\mathcal{V}Z+(\rho,\eta)\, D_{\mathcal{V}[\mathcal{V}X,\mathcal{V}Y]_{(\rho,\eta)TE}}\mathcal{V}Z. \end{cases} \tag{6}$$

Using the above theorem, we obtain the following formulas of Ricci type:

$$\begin{cases} \tilde{Z}^{\alpha}{}_{|\gamma|\beta}-\tilde{Z}^{\alpha}{}_{|\beta|\gamma} & =(\rho,\eta,h)\,\mathbb{R}^{\alpha}_{\theta\ \gamma\beta}\tilde{Z}^{\theta}+\left(L^{\theta}_{\beta\gamma}\circ h\circ\pi\right)\tilde{Z}^{\alpha}{}_{|\theta} \\ & \quad +(\rho,\eta,h)\,\mathbb{T}^{\circ}{}_{\beta\gamma}\tilde{Z}^{\alpha}|_{\circ}+(\rho,\eta,h)\,\mathbb{T}^{\theta}{}_{\beta\gamma}\tilde{Z}^{\alpha}{}_{|\theta}, \\ \tilde{Z}^{\alpha}{}_{|\gamma}|_{\circ}-\tilde{Z}^{\alpha}|_{\circ|\gamma} & =(\rho,\eta,h)\,\mathbb{P}^{\alpha}_{\theta\ \gamma\circ}\tilde{Z}^{\theta}-(\rho,\eta,h)\,\mathbb{P}^{\circ}{}_{\gamma\circ}\tilde{Z}^{\alpha}|_{\circ} \\ & \quad -(\rho,\eta)\,\mathbb{H}^{\circ}_{\circ\gamma}\tilde{Z}^{\alpha}|_{\circ}, \\ \tilde{Z}^{\alpha}|_{\circ}|_{\circ}-\tilde{Z}^{\alpha}|_{\circ}|_{\circ} & =(\rho,\eta,h)\,\mathbb{S}^{\alpha}_{\theta\ \circ\circ}\tilde{Z}^{\theta}+(\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\circ}\tilde{Z}^{\alpha}|_{\circ}, \end{cases} \tag{7}$$

and

$$\begin{cases} Y^{\circ}{}_{|\gamma|\beta}-Y^{\circ}{}_{|\beta|\gamma} & =(\rho,\eta,h)\,\mathbb{R}^{\circ}{}_{\circ\ \gamma\beta}Y^{\circ}+\left(L^{\theta}_{\beta\gamma}\circ h\circ\pi\right)Y^{\circ}{}_{|\theta} \\ & \quad +(\rho,\eta)\,\mathbb{T}^{\circ}{}_{\beta\gamma}Y^{\circ}|_{\circ}+(\rho,\eta,h)\,\mathbb{T}^{\theta}{}_{\beta\gamma}Y^{\circ}{}_{|\theta}, \\ Y^{\circ}{}_{|\gamma}|_{\circ}-Y^{\circ}|_{\circ|\gamma} & =(\rho,\eta,h)\,\mathbb{P}^{\circ}_{\theta\ \gamma\circ}Y^{\theta}-(\rho,\eta,h)\,\mathbb{P}^{\circ}{}_{\gamma\circ}Y^{\circ}|_{\circ} \\ & \quad -(\rho,\eta)\,\mathbb{H}^{\circ}_{\circ\gamma}Y^{\circ}|_{\circ}, \\ Y^{\circ}|_{\circ}|_{\circ}-Y^{\circ}|_{\circ}|_{\circ} & =(\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\ \circ\circ}Y^{\circ}+(\rho,\eta,h)\,\mathbb{S}^{\circ}{}_{\circ\circ}Y^{\circ}|_{\circ}. \end{cases} \tag{8}$$

In particular, if $(\rho,\eta,h)=(Id_{TM},Id_M,id_M)$ and the Lie bracket $[,]_{TM}$ is the usual Lie bracket, then the formulas of Ricci type (7) and (8) reduce to

$$\left\{\begin{array}{ll}\tilde{Z}^i{}_{|k|j}-\tilde{Z}^i{}_{|j|k} & =\mathbb{R}^i{}_{h\ kj}\tilde{Z}^h+\mathbb{T}^\circ{}_{jk}\tilde{Z}^i|_\circ+\mathbb{T}^h{}_{jk}\tilde{Z}^i{}_{|h},\\ \tilde{Z}^i{}_{|k}|_\circ-\tilde{Z}^i|_{\circ|k} & =\mathbb{P}^i{}_{h\ k\circ}\tilde{Z}^h-\mathbb{P}^\circ{}_{k\circ}\tilde{Z}^i|_\circ-\mathbb{H}^\circ{}_{\circ k}\tilde{Z}^i|_\circ,\\ \tilde{Z}^i|_\circ|_\circ-\tilde{Z}^i|_\circ|_\circ & =\mathbb{S}^i{}_{h\ \circ\circ}\tilde{Z}^h+\mathbb{S}^\circ{}_{\circ\circ}\tilde{Z}^i|_\circ,\end{array}\right.$$

and

$$\left\{\begin{array}{ll}Y^\circ{}_{|k|j}-Y^\circ{}_{|j|k} & =\mathbb{R}^\circ{}_{0\ kj}Y^c+\mathbb{T}^\circ{}_{jk}Y^\circ|_\circ+\mathbb{T}^h{}_{jk}Y^\circ{}_{|h},\\ Y^\circ{}_{|k}|_\circ-Y^\circ|_{\circ|k} & =\mathbb{P}^\circ{}_{h\ k\circ}Y^h-\mathbb{P}^\circ{}_{k\circ}Y^\circ|_\circ-\mathbb{H}^\circ{}_{\circ k}Y^\circ|_\circ,\\ Y^\circ|_\circ|_\circ-Y^\circ|_\circ|_\circ & =\mathbb{S}^\circ{}_{\circ\ \circ\circ}Y^\circ+\mathbb{S}^\circ{}_{\circ\circ}Y^\circ|_\circ.\end{array}\right.$$

We consider the following 1-forms associated to distinguished linear (ρ,η)-connection $((\rho,\eta)\,H,(\rho,\eta)\,V)$

$$\begin{aligned}(\rho,\eta)\,\omega^\alpha_\beta&=(\rho,\eta)\,H^\alpha_{\beta\gamma}d\tilde{z}^\gamma+(\rho,\eta)\,V^\alpha_{\beta\circ}\delta\tilde{y}^\circ,\\(\rho,\eta)\,\omega^\circ_\circ&=(\rho,\eta)\,H^\circ_{\circ\gamma}d\tilde{z}^\gamma+(\rho,\eta)\,V^\circ_{\circ\circ}\delta\tilde{y}^\circ.\end{aligned}$$

Moreover, we obtain the following torsion 2-forms

$$\left\{\begin{array}{l}(\rho,\eta,h)\,\mathbb{T}^\alpha=\dfrac{1}{2}\,(\rho,\eta,h)\,\mathbb{T}^\alpha{}_{\beta\gamma}d\tilde{z}^\beta\wedge d\tilde{z}^\gamma+(\rho,\eta,h)\,\mathbb{P}^\alpha{}_{\beta\circ}d\tilde{z}^\beta\wedge\delta\tilde{y}^\circ,\\(\rho,\eta,h)\,\mathbb{T}^\circ=\dfrac{1}{2}\,(\rho,\eta,h)\,\mathbb{T}^\circ{}_{\beta\gamma}d\tilde{z}^\beta\wedge d\tilde{z}^\gamma+(\rho,\eta,h)\,\mathbb{P}^\circ{}_{\beta\circ}d\tilde{z}^\beta\wedge\delta\tilde{y}^\circ\\\qquad\qquad+\dfrac{1}{2}\,(\rho,\eta,h)\,\mathbb{S}^\circ{}_{\circ\circ}\delta\tilde{y}^\circ\wedge\delta\tilde{y}^\circ,\end{array}\right.$$

and

$$\left\{\begin{array}{l}(\rho,\eta,h)\,\mathbb{R}^\alpha{}_\beta=\dfrac{1}{2}\,(\rho,\eta,h)\,\mathbb{R}^\alpha{}_{\beta\ \gamma\theta}d\tilde{z}^\gamma\wedge d\tilde{z}^\theta+(\rho,\eta,h)\,\mathbb{P}^\alpha{}_{\beta\ \gamma\circ}d\tilde{z}^\gamma\wedge\delta\tilde{y}^\circ\\\qquad\qquad+\frac{1}{2}\,(\rho,\eta,h)\,\mathbb{S}^\alpha{}_{\beta\ \circ\circ}\delta\tilde{y}^\circ\wedge\delta\tilde{y}^\circ,\\(\rho,\eta,h)\,\mathbb{R}^\circ{}_\circ=\dfrac{1}{2}\,(\rho,\eta,h)\,\mathbb{R}^\circ{}_{\circ\ \gamma\theta}d\tilde{z}^\gamma\wedge d\tilde{z}^\theta+(\rho,\eta,h)\,\mathbb{P}^\circ{}_{\circ\ \gamma\circ}d\tilde{z}^\gamma\wedge\delta\tilde{y}^\circ\\\qquad\qquad+\frac{1}{2}\,(\rho,\eta,h)\,\mathbb{S}^\circ{}_{\circ\ \circ\circ}\delta\tilde{y}^\circ\wedge\delta\tilde{y}^\circ.\end{array}\right.$$

Theorem 29 *The following identities of Cartan type hold*

$$\begin{aligned}(\rho,\eta,h)\,\mathbb{T}^\alpha&=d^{(\rho,\eta)TE}\,(d\tilde{z}^\alpha)+(\rho,\eta)\,\omega^\alpha_\beta\wedge d\tilde{z}^\beta,\\(\rho,\eta,h)\,\mathbb{T}^\circ&=d^{(\rho,\eta)TE}\,(\delta\tilde{y}^\circ)+(\rho,\eta)\,\omega^\circ_\circ\wedge\delta\tilde{y}^\circ,\end{aligned}\tag{9}$$

and

$$\begin{aligned}(\rho,\eta,h)\,\mathbb{R}^\alpha{}_\beta&=d^{(\rho,\eta)TE}\left((\rho,\eta)\,\omega^\alpha_\beta\right)+(\rho,\eta)\,\omega^\alpha_\gamma\wedge(\rho,\eta)\,\omega^\gamma_\beta,\\(\rho,\eta,h)\,\mathbb{R}^\circ{}_\circ&=d^{(\rho,\eta)TE}\,((\rho,\eta)\,\omega^\circ_\circ)+(\rho,\eta)\,\omega^\circ_\circ\wedge(\rho,\eta)\,\omega^\circ_\circ.\end{aligned}\tag{10}$$

In particular, if $(\rho,\eta,h)=(Id_{TM},Id_M,Id_M)$ and the Lie bracket $[,]_{TM}$ is the usual Lie bracket, then the identities of Cartan type (9) and (10) reduce to

$$\begin{aligned}\mathbb{T}^i &= d^{(Id_{TE},Id_E)TE}\left(d\tilde{z}^i\right)+\omega^i_j\wedge d\tilde{z}^j,\\ \mathbb{T}^\circ &= d^{(Id_{TE},Id_E)TE}\left(\delta\tilde{y}^\circ\right)+\omega^\circ_\circ\wedge\delta\tilde{y}^\circ,\end{aligned}$$

and

$$\begin{aligned}\mathbb{R}^i{}_j &= d^{(Id_{TE},Id_E)TE}\left(\omega^i_j\right)+\omega^i_k\wedge\omega^k_j,\\ \mathbb{R}^\circ{}_\circ &= d^{(Id_{TE},Id_E)TE}\left(\omega^\circ_\circ\right)+\omega^\circ_\circ\wedge\omega^\circ_\circ.\end{aligned}$$

Remark 30 *For any $X,Y,Z\in\Gamma\left((\rho,\eta)TE,(\rho,\eta)\tau_E,E\right)$, the following identities hold*

$$\begin{aligned}\mathcal{V}(\rho,\eta,h)\mathbb{R}(X,Y)\mathcal{H}Z &= 0,\\ \mathcal{H}(\rho,\eta,h)\mathbb{R}(X,Y)\mathcal{V}Z &= 0,\\ \mathcal{V}D_X\left((\rho,\eta,h)\mathbb{R}(Y,Z)\mathcal{H}U\right) &= 0,\\ \mathcal{H}D_X\left((\rho,\eta,h)\mathbb{R}(Y,Z)\mathcal{V}U\right) &= 0,\end{aligned}$$

and

$$(\rho,\eta,h)\mathbb{R}(X,Y)Z=\mathcal{H}(\rho,\eta,h)\mathbb{R}(X,Y)\mathcal{H}Z+\mathcal{V}(\rho,\eta,h)\mathbb{R}(X,Y)\mathcal{V}Z.$$

Using the formulas of Bianchi type, Theorem 29 and Remark 30, we obtain the following

Theorem 31 *The identities of Bianchi type:*

$$\begin{cases}\sum\limits_{cyclic(X,Y,Z)}\{\mathcal{H}(\rho,\eta)D_X\left((\rho,\eta,h)\mathbb{T}(Y,Z)\right)-\mathcal{H}(\rho,\eta,h)\mathbb{R}(X,Y)Z\\ \qquad+\mathcal{H}(\rho,\eta,h)\mathbb{T}\left(\mathcal{H}(\rho,\eta,h)\mathbb{T}(X,Y),Z\right)\\ \qquad+\mathcal{H}(\rho,\eta,h)\mathbb{T}\left(\mathcal{V}(\rho,\eta,h)\mathbb{T}(X,Y),Z\right)\}=0,\\ \sum\limits_{cyclic(X,Y,Z)}\{\mathcal{V}(\rho,\eta)D_X\left((\rho,\eta,h)\mathbb{T}(Y,Z)\right)-\mathcal{V}(\rho,\eta,h)\mathbb{R}(X,Y)Z\\ \qquad+\mathcal{V}(\rho,\eta,h)\mathbb{T}\left(\mathcal{H}(\rho,\eta,h)\mathbb{T}(X,Y),Z\right)\\ \qquad+\mathcal{V}(\rho,\eta,h)\mathbb{T}\left(\mathcal{V}(\rho,\eta,h)\mathbb{T}(X,Y),Z\right)\}=0.\end{cases}$$

and

$$\begin{cases} \sum\limits_{cyclic(X,Y,Z,U)} \{\mathcal{H}(\rho,\eta) D_X((\rho,\eta,h)\mathbb{R}(Y,Z)U) \\ \qquad -\mathcal{H}(\rho,\eta,h)\mathbb{R}(\mathcal{H}(\rho,\eta,h)\mathbb{T}(X,Y),Z)U \\ \qquad -\mathcal{H}(\rho,\eta,h)\mathbb{R}(\mathcal{V}(\rho,\eta,h)\mathbb{T}(X,Y),Z)U\} = 0, \\ \sum\limits_{cyclic(X,Y,Z,U)} \{\mathcal{V}(\rho,\eta) D_X((\rho,\eta,h)\mathbb{R}(Y,Z)U) \\ \qquad -\mathcal{V}(\rho,\eta,h)\mathbb{R}(\mathcal{H}(\rho,\eta,h)\mathbb{T}(X,Y),Z)U \\ \qquad -\mathcal{V}(\rho,\eta,h)\mathbb{R}(\mathcal{V}(\rho,\eta,h)\mathbb{T}(X,Y),Z)U\} = 0, \end{cases}$$

hold for any $X,Y,Z \in \Gamma((\rho,\eta)TE,(\rho,\eta)\tau_E,E)$.

Corollary 32 *Using the following sections* $(\delta_\theta,\delta_\gamma,\delta_\beta)$*, we obtain:*

$$\begin{cases} \sum\limits_{cyclic(\beta,\gamma,\theta)} \Big\{(\rho,\eta,h)\mathbb{T}^{\alpha}{}_{\beta\gamma_{|\theta}} - (\rho,\eta,h)\mathbb{R}^{\alpha}{}_{\beta\ \gamma\theta} \\ \qquad + (\rho,\eta,h)\mathbb{T}^{\lambda}{}_{\gamma\theta}(\rho,\eta,h)\mathbb{T}^{\alpha}{}_{\beta\gamma} + (\rho,\eta,h)\mathbb{T}^{\circ}{}_{\gamma\theta}(\rho,\eta,h)\mathbb{T}^{\alpha}{}_{\beta\circ}\Big\} = 0, \\ \sum\limits_{cyclic(\beta,\gamma,\theta)} \Big\{(\rho,\eta,h)\mathbb{T}^{\circ}{}_{\beta\gamma_{|\theta}} + (\rho,\eta,h)\mathbb{T}^{\alpha}{}_{\gamma\theta}(\rho,\eta,h)\mathbb{P}^{\circ}{}_{\beta\alpha} \\ \qquad + (\rho,\eta,h)\mathbb{P}^{\circ}{}_{\gamma\theta}(\rho,\eta,h)\mathbb{P}^{\circ}{}_{\circ\beta}\Big\} = 0, \end{cases}$$

and

$$\begin{cases} \sum\limits_{cyclic(\beta,\gamma,\theta,\lambda)} \Big\{(\rho,\eta,h)\mathbb{R}^{\alpha}{}_{\beta\ \gamma\theta_{|\lambda}} - (\rho,\eta,h)\mathbb{T}^{\mu}{}_{\theta\lambda}(\rho,\eta,h)\mathbb{R}^{\alpha}{}_{\beta\ \gamma\mu} \\ \qquad - (\rho,\eta,h)\mathbb{T}^{\circ}{}_{\theta\lambda}(\rho,\eta,h)\mathbb{P}^{\alpha}{}_{\beta\ \gamma\circ}\Big\} = 0, \\ \sum\limits_{cyclic(\beta,\gamma,\theta,\lambda)} \Big\{(\rho,\eta,h)\mathbb{R}^{\circ}{}_{\circ\ \gamma\theta_{|\lambda}} - (\rho,\eta,h)\mathbb{T}^{\mu}{}_{\theta\lambda}(\rho,\eta,h)\mathbb{R}^{\circ}{}_{\circ\ \gamma\mu} \\ \qquad - (\rho,\eta,h)\mathbb{T}^{\circ}{}_{\theta\lambda}(\rho,\eta,h)\mathbb{P}^{\circ}{}_{\circ\ \gamma\circ}\Big\} = 0. \end{cases}$$

Note that using another base of sections, we can obtain new identities of Bianchi type necessary in the applications.

6 (Pseudo) generalized Kaluza-Klein G-spaces

Let $(\rho,\eta)\Gamma$ be a (ρ,η)-connection for the Kaluza-Klein bundle (E,π,M) and let $((\rho,\eta)H,(\rho,\eta)V)$ be a distinguished linear (ρ,η)-connection for the Lie algebroid generalized tangent bundle

$$\Big(((\rho,\eta)TE,(\rho,\eta)\tau_E,E),[,]_{(\rho,\eta)TE},(\tilde{\rho},Id_E)\Big).$$

We consider the following tensor field

$$G = g_{\alpha\beta} d\tilde{z}^{\alpha} \otimes d\tilde{z}^{\beta} + g_{\circ\circ} \delta\tilde{y}^{\circ} \otimes \delta\tilde{y}^{\circ} \in \mathcal{T}_{22}^{00}\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right).$$

If G is symmetric and for any point $u_x \in E$, matrice $(g_{\alpha\beta}(u_x))$ is nondegenerate and also $g_{\circ\circ}(u_x)$ is invertible, then G is called *pseudometrical structure*. Moreover, if $(g_{\alpha\beta}(u_x))$ and $g_{\circ\circ}(u_x)$ have constant signature, then G is called *metrical structure*.

If $\alpha, \beta \in 1, \ldots, p$, then for any local fiber chart (U, s_U) of (E, π, M), we consider the real functions

$$\pi^{-1}(U) \quad \xrightarrow{\tilde{g}^{\beta\alpha}} \mathbb{R},$$

and

$$\pi^{-1}(U) \quad \xrightarrow{\tilde{g}^{\circ\circ}} \mathbb{R},$$

such that

$$\left(\tilde{g}^{\beta\alpha}(u_x)\right) = \left(g_{\alpha\beta}(u_x)\right)^{-1},$$

and

$$\tilde{g}^{\circ\circ}(u_x) = g_{\circ\circ}(u_x)^{-1},$$

for any $u_x \in \pi^{-1}(U) \setminus \{0_x\}$.

If around each point $x \in M$ there exists a local vector chart (U, s_U) and a local chart (U, ξ_U) such that $g_{\alpha\beta} \circ s_U^{-1} \circ \left(\xi_U^{-1} \times Id_{\mathbb{R}}\right)(x, y)$ (respectively $g_{\circ\circ} \circ s_U^{-1} \circ \left(\xi_U^{-1} \times Id_{\mathbb{R}}\right)(x, y)$) depends only on x, for any $u_x \in \pi^{-1}(U)$, then we say that the (pseudo)metrical structure G is is a *Riemannian $\mathcal{H}$-(pseudo) metrical structure* (respectively, *Riemannian $\mathcal{V}$-(pseudo)metrical structure*). Moreover, G is called *Riemannian (pseudo) metrical structure* if it is $\mathcal{H}$- and $\mathcal{V}$-(pseudo) metrical structures.

Definition 33 *If there exists a (pseudo) metrical structure*

$$G = g_{\alpha\beta} d\tilde{z}^{\alpha} \otimes d\tilde{z}^{\beta} + g_{\circ\circ} \delta\tilde{y}^{\circ} \otimes \delta\tilde{y}^{\circ}, \tag{11}$$

such that

$$(\rho,\eta)\, D_X G = 0, \ \forall X \in \Gamma\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right), \tag{12}$$

then the Lie algebroid generalized tangent bundle

$$\left(\left((\rho,\eta)\, TE, (\rho,\eta)\, \tau_E, E\right), [,]_{(\rho,\eta)TE}, (\tilde{\rho}, Id_E)\right),$$

is called (pseudo) generalized Kaluza-Klein G-space. Moreover, the local real functions $(\rho,\eta)\Gamma_{\alpha}^{\circ}$ and $(\rho,\eta)R^{\circ}{}_{\alpha\beta}$ are called the electromagnetic potentials and the components of the electromagnetic field, respectively.

One can deduce that the condition (12) is equivalent with the following:

$$g_{\alpha\beta|\gamma} = 0, \quad g_{\circ\circ|\gamma} = 0, \quad g_{\alpha\beta}\,|_{\circ} = 0\,, \ g_{\circ\circ}\,|_{\circ} = 0.$$

If $g_{\alpha\beta|\gamma}$=0 and $g_{\circ\circ|\gamma}$=0, then we will say that the Lie algebroid generalized tangent bundle

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_E,E\right),\left[,\right]_{(\rho,\eta)TE},\left(\tilde{\rho},Id_E\right)\right),$$

is $\mathcal{H}$*-(pseudo) generalized Kaluza-Klein G-space.* Also, if $g_{\alpha\beta}|_{\circ}$=0 and $g_{\circ\circ}|_{\circ}$=0, then we say that the Lie algebroid generalized tangent bundle

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_E,E\right),\left[,\right]_{(\rho,\eta)TE},\left(\tilde{\rho},Id_E\right)\right),$$

is $\mathcal{V}$*-(pseudo) generalized Kaluza-Klein G-space.*

Theorem 34 *If* $\left(\left(\rho,\eta\right)\mathring{H},\left(\rho,\eta\right)\mathring{V}\right)$ *is a distinguished linear* (ρ,η)*-connection for the Lie algebroid generalized tangent bundle*

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_E,E\right),\left[,\right]_{(\rho,\eta)TE},\left(\tilde{\rho},Id_E\right)\right),$$

and G is a (pseudo) metrical structure given by (11), then the real local functions:

$$\begin{aligned}
\left(\rho,\eta\right)H^{\alpha}_{\beta\gamma} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)g_{\varepsilon\beta}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\beta}\right)g_{\varepsilon\gamma}-\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\varepsilon}\right)g_{\beta\gamma}\right.\\
&\left.+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ h\circ\pi-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ h\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ h\circ\pi\right),\\
\left(\rho,\eta\right)H^{\circ}_{\circ\gamma} &= \left(\rho,\eta\right)\mathring{H}^{\circ}_{\circ\gamma}+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\overset{\circ}{\circ\circ}|\gamma},\\
\left(\rho,\eta\right)V^{\alpha}_{\beta\circ} &= \left(\rho,\eta\right)\mathring{V}^{\alpha}_{\beta\circ}+\frac{1}{2}\tilde{g}^{\alpha\varepsilon}g_{\beta\varepsilon|\overset{\circ}{\circ}},\\
\left(\rho,\eta\right)V^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}-\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}\right),
\end{aligned}$$

are the components of a distinguished linear (ρ,η)*-connection such that the Lie algebroid generalized tangent bundle*

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_E,E\right),\left[,\right]_{(\rho,\eta)TE},\left(\tilde{\rho},Id_E\right)\right),$$

becomes (pseudo) generalized Kaluza-Klein G-space.

Corollary 35 *In the particular case of Lie algebroids,* $(\eta,h)=(Id_M,Id_M)$*, then we obtain*

$$\begin{aligned}
\rho H^{\alpha}_{\beta\gamma} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)g_{\varepsilon\beta}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\beta}\right)g_{\varepsilon\gamma}-\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\varepsilon}\right)g_{\beta\gamma}\right.\\
&\left.+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ\pi-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ\pi\right),\\
\rho H^{\circ}_{\circ\gamma} &= \rho\mathring{H}^{\circ}_{\circ\gamma}+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\overset{\circ}{\circ\circ}|\gamma},\\
\rho V^{\alpha}_{\beta\circ} &= \rho\mathring{V}^{\alpha}_{\beta\circ}+\frac{1}{2}\tilde{g}^{\alpha\varepsilon}g_{\beta\varepsilon|\overset{\circ}{\circ}},\\
\rho V^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}-\Gamma\left(\tilde{\rho},Id_E\right)\left(\dot{\tilde{\partial}}_{\circ}\right)g_{\circ\circ}\right)
\end{aligned}$$

In the classical case, $(\rho,\eta,h)=(Id_{TM},Id_M,Id_M)$, we obtain

$$\begin{aligned}
H^i_{jk} &= \frac{1}{2}\tilde{g}^{ih}\left(\delta_k g_{hj}+\delta_j g_{hk}-\delta_h g_{jk}\right),\\
H^{\circ}_{\circ k} &= \mathring{H}^{\circ}_{\circ k}+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\circ\circ|k},\\
V^i_{j\circ} &= \mathring{V}^i_{j\circ}+\frac{1}{2}\tilde{g}^{ih}g_{jh|\circ},\\
V^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\left(\dot{\partial}_{\circ}g_{\circ\circ}+\dot{\partial}_{\circ}g_{\circ\circ}-\dot{\partial}_{\circ}g_{\circ\circ}\right).
\end{aligned}$$

Theorem 36 *If the distinguished linear (ρ,η)-connection $\left((\rho,\eta)\mathring{H},(\rho,\eta)\mathring{V}\right)$ coincides with the Berwald linear (ρ,η)-connection in Theorem 34, then the local real functions:*

$$\begin{aligned}
(\rho,\eta)\overset{c}{H}{}^{\alpha}_{\beta\gamma} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)g_{\varepsilon\beta}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\beta}\right)g_{\varepsilon\gamma}\right.\\
&\quad -\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\varepsilon}\right)g_{\beta\gamma}+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ h\circ\pi\\
&\quad \left.-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ h\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ h\circ\pi\right),\\
(\rho,\eta)\overset{c}{H}{}^{\circ}_{\circ\gamma} &= \dot{\partial}_{\circ}\left((\rho,\eta)\Gamma^{\circ}_{\gamma}\right)+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\circ\circ|\gamma},\\
(\rho,\eta)\overset{c}{V}{}^{\alpha}_{\beta\circ} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\dot{\partial}_{\circ}g_{\beta\varepsilon},\\
(\rho,\eta)\overset{c}{V}{}^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\dot{\partial}_{\circ}g_{\circ\circ},
\end{aligned}$$

are the components of a distinguished linear (ρ,η)-connection such that the Lie algebroid generalized tangent bundle

$$\left(\left((\rho,\eta)TE,(\rho,\eta)\tau_E,E\right),[,]_{(\rho,\eta)TE},\left(\tilde{\rho},Id_E\right)\right),$$

becomes (pseudo) generalized Kaluza-Klein G-space.

Moreover, if the (pseudo) metrical structure G is $\mathcal{H}$- and $\mathcal{V}$-Riemannian, then the local real functions:

$$\begin{aligned}
(\rho,\eta)\overset{c}{H}{}^{\alpha}_{\beta\gamma} &= \tfrac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\rho^k_{\gamma}\circ h\circ\pi\tfrac{\partial g_{\varepsilon\beta}}{\partial x^k}+\rho^j_{\beta}\circ h\circ\pi\tfrac{\partial g_{\varepsilon\gamma}}{\partial x^j}-\rho^e_{\varepsilon}\circ h\circ\pi\tfrac{\partial g_{\beta\gamma}}{\partial x^e}+\right.\\
&\quad \left.+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ h\circ\pi-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ h\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ h\circ\pi\right),\\
(\rho,\eta)\overset{c}{H}{}^{\circ}_{\circ\gamma} &= \dot{\partial}_{\circ}\left((\rho,\eta)\Gamma^{\circ}_{\gamma}\right)+\tfrac{1}{2}\tilde{g}^{\circ\circ}\left(\rho^i_{\gamma}\circ h\circ\pi\tfrac{\partial g_{\circ\circ}}{\partial x^i}-2\dot{\partial}_{\circ}\left((\rho,\eta)\Gamma^{\circ}_{\gamma}\right)\right),\\
(\rho,\eta)\overset{c}{V}{}^{\alpha}_{\beta\circ} &= 0,\\
(\rho,\eta)\overset{c}{V}{}^{\circ}_{\circ\circ} &= 0.
\end{aligned}$$

are the components of a distinguished linear (ρ,η)-connection such that the Lie algebroid generalized tangent bundle becomes (pseudo) generalized Kaluza-Klein G-space.

Corollary 37 *In the particular case of Lie algebroids, $(\eta,h)=(Id_M,Id_M)$, then we obtain*

$$\begin{aligned}
\overset{c}{\rho H}{}^{\alpha}_{\beta\gamma} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\gamma}\right)g_{\varepsilon\beta}+\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\beta}\right)g_{\varepsilon\gamma}\right.\\
&\quad\left.-\Gamma\left(\tilde{\rho},Id_E\right)\left(\tilde{\delta}_{\varepsilon}\right)g_{\beta\gamma}+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ\pi-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ\pi\right),\\
\overset{c}{\rho H}{}^{\circ}_{\circ\gamma} &= \dot{\partial}_{\circ}(\rho\Gamma^{\circ}_{\gamma})+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\circ\circ\overset{\circ}{|}\gamma},\\
\overset{c}{\rho V}{}^{\alpha}_{\beta\circ} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\dot{\partial}_{\circ}g_{\beta\varepsilon},\\
\overset{c}{\rho V}{}^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\dot{\partial}_{\circ}g_{\circ\circ}.
\end{aligned}$$

If the (pseudo) metrical structure G is $\mathcal{H}$-and $\mathcal{V}$-Riemannian, then

$$\begin{aligned}
\overset{c}{\rho H}{}^{\alpha}_{\beta\gamma} &= \frac{1}{2}\tilde{g}^{\alpha\varepsilon}\left(\rho^{k}_{\gamma}\circ\pi\frac{\partial g_{\varepsilon\beta}}{\partial x^{k}}+\rho^{j}_{\beta}\circ\pi\frac{\partial g_{\varepsilon\gamma}}{\partial x^{j}}-\rho^{e}_{\varepsilon}\circ\pi\frac{\partial g_{\beta\gamma}}{\partial x^{e}}+\right.\\
&\quad\left.+g_{\theta\varepsilon}L^{\theta}_{\gamma\beta}\circ\pi-g_{\beta\theta}L^{\theta}_{\gamma\varepsilon}\circ\pi-g_{\theta\gamma}L^{\theta}_{\beta\varepsilon}\circ\pi\right),\\
\overset{c}{\rho H}{}^{\circ}_{\circ\gamma} &= \dot{\partial}_{\circ}(\rho\Gamma^{\circ}_{\gamma})+\tfrac{1}{2}\tilde{g}^{\circ\circ}\rho^{i}_{\gamma}\circ\pi\tfrac{\partial g_{\circ\circ}}{\partial x^{i}},\\
\overset{c}{\rho V}{}^{\alpha}_{\beta\circ} &= 0,\ \overset{c}{\rho V}{}^{0}_{00}=0.
\end{aligned}$$

In the classical case, $(\rho,\eta,h)=(Id_{TM},Id_M,Id_M)$, we obtain

$$\begin{aligned}
\overset{c}{H}{}^{i}_{jk} &= \frac{1}{2}\tilde{g}^{ih}\left(\delta_k g_{hj}+\delta_j g_{hk}-\delta_h g_{jk}\right),\\
\overset{c}{H}{}^{\circ}_{\circ k} &= \dot{\partial}_{\circ}\Gamma^{\circ}_{k}+\frac{1}{2}\tilde{g}^{\circ\circ}g_{\circ\circ\overset{\circ}{|}k},\\
\overset{c}{V}{}^{i}_{j\circ} &= \frac{1}{2}\tilde{g}^{ih}\dot{\partial}_{\circ}g_{jh},\\
\overset{c}{V}{}^{\circ}_{\circ\circ} &= \frac{1}{2}\tilde{g}^{\circ\circ}\dot{\partial}_{\circ}g_{\circ\circ}.
\end{aligned}$$

If the (pseudo)metrical structure G is $\mathcal{H}$- and $\mathcal{V}$-Riemannian, then

$$\begin{aligned}
\overset{c}{H}{}^{i}_{jk} &= \frac{1}{2}\tilde{g}^{ih}\left(\frac{\partial g_{hj}}{\partial x^{k}}+\frac{\partial g_{hk}}{\partial x^{j}}-\frac{\partial g_{jk}}{\partial x^{h}}\right),\\
\overset{c}{H}{}^{\circ}_{\circ k} &= \dot{\partial}_{\circ}\Gamma^{\circ}_{k}+\tfrac{1}{2}\tilde{g}^{\circ\circ}\tfrac{\partial g_{\circ\circ}}{\partial x^{k}},\\
\overset{c}{V}{}^{i}_{j\circ} &= 0,\ \overset{c}{V}{}^{\circ}_{\circ\circ}=0.
\end{aligned}$$

Let

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_{E},E\right),\left[,\right]_{\left(\rho,\eta\right)TE},\left(\tilde{\rho},Id_{E}\right)\right),$$

be a (pseudo) generalized Kaluza-Klein G-space.

Definition 38 *If $\left(\rho,\eta,h\right)\mathbb{R}_{\ \alpha\ \beta}$ and $\left(\rho,\eta,h\right)\mathbb{S}_{\circ\ \circ}$ are the components of Ricci tensor associated to distinguished linear (ρ,η)-connection*

$$\left(\left(\rho,\eta\right)H,\left(\rho,\eta\right)V\right),$$

then the scalar

$$\left(\rho,\eta,h\right)\mathbb{R}=\left(\rho,\eta,h\right)\mathbb{R}_{\ \alpha\ \beta}\tilde{g}^{\alpha\beta}+\left(\rho,\eta,h\right)\mathbb{S}_{\ \circ\ \circ}\tilde{g}^{\circ\circ},$$

is called the scalar curvature of the distinguished linear (ρ,η)-connection $((\rho,\eta)H,(\rho,\eta)V)$.

Definition 39 *The tensor field*

$$\begin{aligned}\left(\rho,\eta,h\right)\mathbb{T} &=\left(\rho,\eta,h\right)\mathbb{T}_{\ \alpha\ \beta}d\tilde{z}^{\alpha}\otimes d\tilde{z}^{\beta}+\left(\rho,\eta,h\right)\mathbb{T}_{\ \alpha\ \circ}d\tilde{z}^{\alpha}\otimes\delta\tilde{y}^{\circ}\\ &+\left(\rho,\eta,h\right)\mathbb{T}_{\ \circ\ \beta}\delta\tilde{y}^{\circ}\otimes d\tilde{z}^{\alpha}+\left(\rho,\eta,h\right)\mathbb{T}_{\ \circ\ \circ}\delta\tilde{y}^{\circ}\otimes\delta\tilde{y}^{\circ},\end{aligned}$$

such that its components satisfy the following conditions:

$$\begin{aligned}\varkappa\left(\rho,\eta,h\right)\mathbb{T}_{\ \alpha\ \beta} &=\left(\rho,\eta,h\right)\mathbb{R}_{\ \alpha\ \beta}-\frac{1}{2}\left(\rho,\eta,h\right)\mathbb{R}\cdot g_{\alpha\beta},\\ -\varkappa\left(\rho,\eta,h\right)\mathbb{T}_{\ \alpha\ \circ} &=\left(\rho,\eta,h\right)\mathbb{P}_{\ \alpha\ \circ},\\ \varkappa\left(\rho,\eta,h\right)\mathbb{T}_{\ \circ\ \beta} &=\left(\rho,\eta,h\right)\mathbb{P}_{\ \circ\ \beta},\\ \varkappa\left(\rho,\eta,h\right)\mathbb{T}_{\ \circ\ \circ} &=\left(\rho,\eta,h\right)\mathbb{S}_{\ \circ\ \circ}-\frac{1}{2}\left(\rho,\eta,h\right)\mathbb{R}\cdot g_{\circ\circ},\end{aligned}\tag{13}$$

where $\varkappa$ is a constant, is called the energy-momentum tensor field associated to the (pseudo) generalized Kluza-Klein G-space

$$\left(\left(\left(\rho,\eta\right)TE,\left(\rho,\eta\right)\tau_{E},E\right),\left[,\right]_{\left(\rho,\eta\right)TE},\left(\tilde{\rho},Id_{E}\right)\right).$$

Also, the equations (13) are called the Einstein equations associated to the (pseudo) generalized Kluza-Klein G-space.

Formally, the Einstein equations can be written

$$\mathbf{Ric}\left(\left(\rho,\eta\right)H,\left(\rho,\eta\right)V\right)-\frac{1}{2}\left(\rho,\eta,h\right)\mathbb{R}\cdot G=\varkappa\cdot\left(\rho,\eta,h\right)\mathbb{T}.$$

Acknowledgment

The first author would like to give his warmest thanks to Radinesti-Gorj Cultural Sciantifique Society for financial support.

References

[1] C. M. Arcuş, *The generalized Lie algebroids and their applications*, arXiv:1007.1541v2, (2010), 206 pages.

[2] C. M. Arcuş, *Generalized Lie algebroids and connections over pair of diffeomorphic manifolds*, J. Gen. Lie Theory Appl., **7** (2013), 32 pages.

[3] C. M. Arcuş, *Metrizability of the Liealgebroid generalized tangent bundle and (generalized) Lagrange (ρ, η)-spaces*, arXiv:1109.1242v3, (2013), 33 pages.

[4] C. M. Arcuş, *Mechanical systems in the generalized Lie algebroids framework*, Int. J. Geom. Methods Mod. Phys., **11** (2014), 40 pages.

[5] A. Bejancu, *A new point of view on general Kaluza-Klein theories*, Prog. Theor. Phys., **128** (2012), 541-585.

[6] A. Bejancu, *4D Equations of motion and fifth force in a 5D general Kaluza-Klein space*, Gen Relativ Gravit, **45** (2013), 2273-2290.

[7] A. Bejancu, *4D Einstein equations in a 5D general Kaluza-Klein space*, Gen Relativ Gravit, **45** (2013), 2291-2308.

[8] A. Bejancu, *On higher dimensional Kaluza-Klein theories*, Advances in High Energy Physics, (2013), 12 pages.

[9] S. Chatterjee, *Homogeneous cosmological model for a Kaluza-Klein-type metric*, Gen. Relat. Gravit., **18(10)** (1986), 1073-1077.

[10] A. Einstein and P. Bergmann, On a generalization of Kaluza's theory of electricty, Ann. of Math., **39 (3)** (1938), 683-701.

[11] R. L. Fernandes, *Lie Algebroids, Holonomy and Characteristic Classes*, Advances in Mathematics **170** (2002), 119-179.

[12] T. Kaluza, *Zum Unitäts problem in der Physik*, Sitzungsber. Preuss. Akad. Wiss., (1921), 966-972.

[13] M. d. León, J. C. Marrero and E. Martínez, *Lagrangian submanifolds and dynamics on Lie algebroids*, J. Phys. A: Math. Gen., **38** (2005), 241-308.

[14] O. Klein, *Quantentheorie und fünfdimensionale Relativitätstheorie*, Zeitschrift für Physik A, **37(12)** (1926), 895-906.

[15] H. Liu and P. S. Wesson, *Cosmological solutions and their effective properties of matter in Kaluza-Klein theory*, Gen. Relat. Gravit., **30 (3)** (1998), 509-514.

[16] J. M. Overduin and P. S. Wesson, *Kaluza-Klein gravity*, Phys. Rep., **283** (1997), 303-378.

[17] J. Ponce de Leon, *Cosmological models in a Kaluza-Klein theory with variable rest mass*, Gen. Relativ. Gravit., **20 (6)** (1988), 539-550.

[18] J. Ponce de Leon and P. S. Wesson, *Exact solutions and the effective equation of state in KaluzaKlein theory*, J. Math. Phys., **34** (1993), 4080-4092.

[19] L. Popescu, *Geometrical structures on Lie algebroids*, Publ. Math. Debrecen, **72 (1-2)** (2008), 95-109.

[20] L. Popescu, *A note on Poisson-Lie algebroids*, J. Geom. Symmetry Phys., **12** (2008), 63-73.

[21] S. Vacaru, *Einstein-Cartan algebroids and black holes in solitonic backgrounds*, arXiv:gr-qc/ 0501057, (2005), 35 pages.

[22] S. Vacaru, *Clifford algebroids and nonholonomic spinor deformations of Taub-NUT spacetimes*, arXiv:hep-th/0502145, (2005), 30 pages.

[23] S. Vacaru, *Nonholonomic deformations of disk solutions and algebroid symmetries in Einstein and extra dimension gravity*, arXiv:gr-qc/0504095, (2005), 66 pages.

[24] S. Vacaru, *Clifford-Finsler algebroids and nonholonomic Einstein-Dirac structures*, J. Math. Phys., **47** (2006), 20 pages.

[25] S. Vacaru, *Nonholonomic algebroids, Finsler geometry, and Lagrange-Hamilton spaces*, Mathematical Sciences, **6 (18)** (2012), 33 pages.

[26] S. Vacaru, *Almost Kaehler Ricci Flows and Einstein and Lagrange-Finsler Structures on Lie Algebroids*, arXiv:1306.2813, (2013), 40 pages.

[27] P. S. Wesson, Space-Time-Matter; Modern Kaluza-Klein Theory, World Scientific, Singapore (1999).

Constantin M Arcuş
Secondary School "Cornelius Radu"
Radinesti Village, 217196
Gorj County, Romania
Email: c_arcus@radinesti.ro

Esmail Peyghan
Department of Mathematics, Faculty of Science
Arak University
Arak 38156-8-8349, Iran
Email: e-peyghan@araku.ac.ir